

Influence of dietary protein / lipid ratio and fish oil substitution on fatty acid composition and metabolism of Atlantic salmon (*Salmo salar*) reared at summer water temperatures

**Vasileios Karalazos<sup>1</sup>, Eldar Å. Bendiksen<sup>2</sup>, James R. Dick<sup>1</sup>, Douglas R. Tocher<sup>1</sup>, John Gordon Bell<sup>1\*</sup>**

<sup>1</sup>*Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, Scotland, UK*

<sup>2</sup>*BioMar AS, Nordre Gate 11, N-7484, Trondheim, Norway*

*\*Corresponding author: Professor John Gordon Bell, Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, Scotland, UK, tel.: +44-1786-467993, fax.: +44-1786-472133, email: g.j.bell@stir.ac.uk*

*Running title: Protein level and oil source in salmon diets*

*Keywords: Atlantic salmon; Rapeseed oil; Polyunsaturated fatty acids (PUFA);  $\beta$ -oxidation*

*Abbreviations: DE: digestible energy; DP digestible protein; FCR: feed conversion ratio; FM: fish meals; FO: fish oil; HUFA: highly unsaturated fatty acid; PPV: protein productive value; RO: rapeseed oil; SGR: specific growth rate; TGC: thermal growth coefficient; VO: vegetable oil*

## **Abstract**

A factorial, two-way, experimental design was used for this 10-week nutritional trial, aiming to elucidate the interactive effects of decreasing dietary protein/lipid level and substitution of fish oil (FO) with rapeseed oil (RO) on tissue fatty acid (FA) composition and metabolism of large Atlantic salmon (*Salmo salar* L.) reared at summer water temperatures (11.6 °C). The six experimental diets were isoenergetic and formulated to include either fish oil (FO) or rapeseed oil (RO - 60% of the added oil) at three dietary protein/lipid levels, specifically 350 g/kg / 350 g/kg,

330 g/kg / 360 g/kg and 290 g/kg / 380 g/kg of protein/lipid. Final weight, SGR and TGC were positively affected by the dietary RO inclusion at the expense of FO, while no significant effects were seen on growth due to the decreasing protein level. The oil source had a significant effect on muscle and liver FA composition. However, the changes in muscle and liver FA indicate selective utilization or retention of individual FA and moderate reductions in tissue EPA and DHA. Pyloric caeca phospholipid FA composition was significantly affected by the two factors and, in some cases, significant interactions were also revealed. Liver and red muscle  $\beta$ -oxidation capacities were significantly increased due to RO inclusion, while an interactive effect of the protein level and the oil source was shown for the white muscle  $\beta$ -oxidation capacity. The results could explain, at least partially, the better performance that was shown for the RO groups and the enhanced protein sparing effect.

## Introduction

In recent years, one of the main research topics in aquaculture nutrition has been the replacement of fish meals (FM) and fish oils (FO) in the diets for fish exploiting alternative sources for protein and lipid, respectively. The use of FM and FO in fish nutrition, especially in the intensive culture of carnivorous species such as Atlantic salmon has been common practice for years and, to a great extent, still is today. This is because they constitute excellent sources of essential amino acids and fatty acids (FA), and especially of highly unsaturated fatty acids (HUFA)<sup>(1-3)</sup>. However, the need for reduction in the consumption of these commodities in aqua feeds is required for a number of reasons, including environmental and economic concerns, including sustainability issues, price increases etc<sup>(4-8)</sup>. In addition, there are issues regarding the quality of the final product, as there is a potential risk of contamination of FM and FO with organic pollutants<sup>(9-13)</sup>.

Numerous commodities have been successfully tested as alternatives to FM and FO. In the case of FO replacement, vegetable oils (VO), including rapeseed oil (RO), have been included, either as single replacements or as part of VO blends. The effects of such dietary alterations have been recently reviewed<sup>(13-15)</sup> and can be summarized as causing no detrimental effects on growth and feed utilization<sup>(16-22)</sup> or notably, in some cases enhancing growth<sup>(21,23,24)</sup>. However, the dietary inclusion of RO at expense of FO leads to significant changes in tissue FA compositions, which reflect the FA compositions of the diets, and FA metabolism, including  $\beta$ -oxidation. It should be noted that RO's content of 18:2n-6 and 18:3n-3 is moderate and at a ratio of 2:1 and thus, should result only in modest deposition of these FA in fish tissues and, perhaps, enhance the endogenous conversion of 18:3n-3 to 20:5n-3 and 22:6n-3. Also, RO contains high levels of monounsaturated

FA, especially 18:1n-9, which are preferred substrates for energy production by Atlantic salmon and hence, growth rates of the fish should not be compromised<sup>(16,25-27)</sup>. Alterations of fish tissue FA profile are of importance given the role of HUFA and especially EPA and DHA, to human health and the increasing demand of consumers for nutritious and health promoting products<sup>(13)</sup>. Fish are unique sources of these FA and hence, their nutritional characteristics should not be compromised.

However, there are issues regarding FM and FO replacement in fish diets which remain unclear. For instance, the authors of the afore mentioned reviews have pointed out that the simultaneous reduction of both FM and FO could be challenging due to the expected reduction in essential FA and amino acids.

In those terms, the investigation of the replacement of FO with a vegetable oil like RO, with a concurrent reduction of FM is of considerable interest. The use of energy dense diets can be a potentially useful approach towards the reduction of FM, as such diet formulations, require less protein by exploiting lipids for energy production and hence, can be potentially advantageous for the nutrition of carnivorous species, like Atlantic salmon<sup>(28,29)</sup>. Research findings agree that these diets are performing well in terms of the fish growth and feed utilization, while a sparing effect on protein by increased dietary oil has also been shown<sup>(23,24,30-34)</sup>. However, in most of these studies relatively high dietary protein/lipid levels were used and, hence, how much protein reduction can occur, without performance loss, still needs to be elucidated. Moreover, the dietary protein/lipid content and the FO replacement with RO, affect tissue FA composition and FA metabolism, including catabolism via  $\beta$ -oxidation which influences fish growth, while other interactions may also occur<sup>(14,23,24,35)</sup>.

In an earlier trial we investigated the interactive effects of the dietary protein/lipid ratio and the inclusion of RO at the expense of FO in the diets of salmon in cold (winter) water temperatures<sup>(24)</sup>. The results of that study showed a positive effect on fish growth, as well as a protein sparing effect but also changes in tissues FA composition. Given that, water temperature is a key-factor in fish nutrition and FA retention and metabolism<sup>(36)</sup>, the impact of even lower dietary protein/lipid ratios and different lipid sources, in Atlantic salmon reared at high (summer) water temperatures is of considerable interest. Moreover, the positive effects of RO inclusion on growth have been associated with changes in the digestibility of FA. The uptake and digestibility of lipids and ultimately the utilization of diets could be affected by the intestine phospholipid FA composition and the consequent alterations occurring due to dietary FA changes, especially when high lipid diets are used.

Hence, the aim of this study was to elucidate the interactive effects of FO replacement with RO at various dietary protein/lipid ratios on tissue FA compositions and metabolism, including FA

$\beta$ -oxidation and phospholipid FA compositions of pyloric caeca, of large Atlantic salmon at high water temperatures.

## **Materials and methods**

### *Fish and facilities*

The 10 week feeding trial was carried out at the Fjord Research Station AS (Helgeland, Dønna, Norway, 66°N) using Atlantic salmon (*Salmo salar*) of initial mean weight of 2053g. Eighteen sea cages of 125 m<sup>3</sup> (5x5x5m) were used with approximately 93 fish randomly distributed in each cage. Prior to the experiment the fish were acclimatised to the trial cages for 63 days at 12°C, being fed a commercial diet from BioMar AS (9 mm; 360 g/kg protein and 350 g/kg fat). During the experimental period the average water temperature was 11.6 ± 1.1 °C and the salinity 32.5 ± 0.4 g L<sup>-1</sup>, while the fish were subjected to natural photoperiod. Mortalities were recorded and dead fish were removed daily. Experimental procedures complied with the Norwegian code of practice for the care and use of animals for scientific purposes. There are no aspects of this trial that would cause aggravated or unnecessary harm or stress to the fish involved.

### *Experimental diets and feeding*

Following a factorial (two-way 3x2) experimental design, six diets were formulated at three different dietary protein/fat levels and two different levels of FO substitution. Regarding the protein/lipid level the diets contained 350 g/kg / 350 g/kg (high protein - HP), 330 g/kg / 360 g/kg (medium protein - MP) and 290 g/kg / 380 g/kg (low protein - LP). For the oil source, FO or RO were used within each dietary protein/fat level, where crude RO comprised 60% of the total added oil in the RO diets, the remainder being FO. All diets were isoenergetic (gross energy, GE 25 kJ g<sup>-1</sup>). The diets were formulated to meet all known nutritional requirements of salmonid fish<sup>(3)</sup> and were produced as practical-type extruded pellets (9 mm) at the BioMar TechCentre (Brande, DK). The ingredients, proximate and fatty acid compositions of the experimental diets are shown in Tables 1 and 2, respectively. Feeding was carried out by hand to satiation on a daily basis, including two daily meals with a minimum of 4 hours between them. A lift-up system was used to collect uneaten feed, which was recorded for accurate calculations of feed intake and FCR.

### *Sampling procedure*

At the start of the trial the fish were bulk weighed. At the end (10th week) the fish were individually weighed after being anaesthetized in MS-222 (metacain, 8mg/L). At the end of the trial another three fish per cage were sampled at random for lipid and fatty acid composition of muscle, liver and pyloric caeca phospholipids and for  $\beta$ -oxidation determination. Fish were killed with a

sharp blow to the head and samples of liver and pyloric caeca were dissected and immediately frozen in liquid nitrogen. Muscle samples, representative of the edible portion, were obtained by cutting a steak between the dorsal and ventral fins (NQC), which were then skinned, de-boned and homogenized. For  $\beta$ -oxidation determination samples of liver, red and white muscle were taken separately and immediately frozen in liquid nitrogen. Samples from the six experimental diets were also taken to determine proximate and FA compositions. All samples were kept at -20 °C until further analysis.

#### *Lipid extraction and fatty acid analyses*

Total lipids of tissues and diet samples were extracted by homogenization in 20 volumes of chloroform/methanol (2 : 1, v/v) containing butylated hydroxytoluene (0.01% w/w, BHT) as antioxidant<sup>(37)</sup>. Fatty acid methyl esters (FAME) were prepared from total lipid by acid-catalysed transesterification using 2 mL of 1% H<sub>2</sub>SO<sub>4</sub> in methanol plus 1 mL toluene, as described by Christie<sup>(38)</sup>, and FAME extraction and purification as described by Tocher & Harvie<sup>(39)</sup>. FAME were separated and quantified by gas-liquid chromatography (Carlo Erba Vega 8160, Milan, Italy) using a 30 m x 0.32 mm capillary column (CP wax 52CB; Chrompak Ltd., London, UK). The carrier gas was hydrogen and the temperature programming used was from 50 to 150 °C at 40 °C min<sup>-1</sup> and then to 225 °C at 2 °C min<sup>-1</sup>. Individual methyl esters were identified by comparison with known standards and by reference to published data<sup>(40)</sup>.

In the case of phospholipid FA composition of pyloric caecae, a phospholipid (PL) fraction was prepared from 0.5 mg of total lipid applied to a 20 x 20 cm silica gel 60 TLC plate (VWR, Lutterworth, England) and developed in iso-hexane/diethyl ether/acetic acid (80:20:1 v/v/v) and dried for a few minutes at room temperature. The plate was sprayed lightly with 2, 7, dichlorofluorescein (0.1% w/v) in 97% methanol (v/v) and the PL bands on the origin scraped from the plate and placed in a 15 ml test tube. FAME were prepared by acid-catalysed transesterification in 2 ml of 1% H<sub>2</sub>SO<sub>4</sub> in methanol at 50°C overnight<sup>(38)</sup>. The samples were neutralised with 2.5 ml of 2% KHCO<sub>3</sub> and extracted with 5 ml isohexane/diethyl ether (1:1 v/v) + BHT. The samples were then re-extracted with 5 ml isohexane/diethyl ether (1:1) and the combined extracts dried and dissolved in 0.3 ml of isohexane prior to fatty acid analysis.

#### *Peroxisomal $\beta$ -oxidation capacity*

Liver and red and white muscle were weighed and homogenized in 20% (w/v) ice-cold buffered sucrose solution containing 0.25M sucrose, 0.04M potassium phosphate buffer (pH 7.4), 0.15M KCl, 40mM KF and 1mM N-acetyl cysteine. The resulting total homogenates were then centrifuged at 1880 × g for 10 min at 2°C. The resulting post-nuclear fractions were collected, and

portions were used immediately to determine  $\beta$ -oxidation capacity. The latter was determined as acid-soluble products using radiolabelled [1- $^{14}\text{C}$ ]-palmitoyl-CoA as a substrate as described by Frøyland *et al*<sup>(41)</sup>.

Briefly, 250  $\mu\text{L}$  of assay medium were added to 2ml eppendorf tubes. Then 10  $\mu\text{L}$  of [1- $^{14}\text{C}$ ]-palmitoyl-CoA substrate (0.1  $\mu\text{Ci}/100 \mu\text{M}$ ) were added to each tube. The samples were pre-incubated at room temperature for 2 min. The reaction was started by the addition of homogenate (30-50  $\mu\text{L}$  for liver and red muscle and 300-500  $\mu\text{L}$  for white muscle, homogenized in 20% (w/v) ice-cold buffered sucrose solution as described above) and the reaction continued for 10 min. The reaction was stopped by addition of 150  $\mu\text{L}$  of 1.5 M KOH. Then 25  $\mu\text{L}$  FAF-BSA (100mg/ml) were added, the tubes were vortexed and 500  $\mu\text{L}$  of ice-cold 4M  $\text{HClO}_4$  (perchloric acid) were added. The tubes were centrifuged at 1880 x g for 10 min. Aliquots of 500  $\mu\text{L}$  were placed in scintillation vials, 2.5 ml of scintillant added and the radioactivity determined in a scintillation counter. The protein content of the samples was determined according to the Lowry method<sup>(42)</sup>.

#### *Calculations and statistical analysis*

The following formulae were applied to the data:

Feed Conversion Ratio (FCR) = feed intake (g) x wet weight gain<sup>-1</sup> (g)

Specific Growth Rate (SGR, %/day) =  $100 \times [\ln W_1 - \ln W_0] \times (\text{days})^{-1}$

Thermal Growth Coefficient (TGC, x 1000) =  $1000 \times [(W_1)^{1/3} - (W_0)^{1/3}] \times (\text{days} \times \text{Temp. } ^\circ\text{C})^{-1}$

Protein productive value (PPV, g protein gain x g protein ingested<sup>-1</sup>) =  $[(P_1 W_1 - P_0 W_0) \times (P_F \times \text{cumulative feed intake})^{-1}]$

where  $W_0$  and  $W_1$  are the initial and final fish mean weights in grams,  $P_0$  and  $P_1$  are the initial and final protein concentrations of the fish,  $P_F$  is the protein concentration of the feed on a dry matter basis, and cumulative feed intake was determined in grams on a dry matter basis.

Factorial (two-way) ANOVA was used to analyse the effects of the protein/fat ratio (protein level), dietary RO inclusion (oil source) and their interactions on FA composition of tissues and  $\beta$ -oxidation. When the interaction of the two factors was significant, multiple comparison testing was performed for both factors to investigate the simple main effects, that is the main effect of one factor at a given level of the other, while the main effects were not taken into account<sup>(43)</sup>. Data which were identified as non-homogeneous (Levene's test) were subjected to square root, log or arcsin transformation before analysis. Differences were regarded as significant when  $P < 0.05$ <sup>(43)</sup>. All the data are presented as means  $\pm$  SD ( $n = 3$ ) and all statistical analyses were performed using SPSS 14.0 (SPSS Inc, 2005). The graphs were created using Prism 4 (Graphpad Software Inc., San Diego, USA).

## Results

### *Diet proximate and fatty acid composition*

The analysed protein / lipid content of the diets, was 349 g/kg / 350 g/kg, 333 g/kg / 358 g/kg, 293 g/kg / 384 g/kg for HP, MP and LP, respectively (Table 1). The Digestible Protein / Digestible Energy (DP/DE) ratio was 14.5, 13.5 and 12.3 for the HP, MP and LP diets, respectively; DP and DE were calculated using the ADC values for protein and energy found in this trial (Karalazos et al., 2010 Aquaculture, submitted for publication). The diets contained either 100% FO or a blend of 40% FO and 60% RO with a consequential effect on the total lipid FA profiles (Table 2). Briefly, the FO diets contained approximately 36% total saturated FA, largely 16:0 (approx. 20%), except for the HP-FO diet which had a slightly higher total saturated FA content (40.4%). The total monoenes were 26%, predominantly 18:1n-9 and 16:1n-7. The total n-6 PUFA were low (4.5%), half of which was 18:2n-6. Lastly, the total n-3 PUFA were as high as 32%, mainly as EPA and DHA (18.5% and 8%, respectively). The 60% inclusion of RO resulted in the reduction of 16:0, and consequently of the total saturated FA, by half, compared to the FO diets. The total monoenes almost doubled, largely due to the high amount of 18:1n-9 (38.5%). The total n-6 PUFA in the RO diets increased 3-fold, up to 14%, mainly as 18:2n-6 (13% of total FA). EPA decreased by 70% and DHA by more than half (values were 6% and 3%, respectively) and hence the total n-3 PUFA were reduced by half (17%). The n-3/n-6 PUFA ratio was 7 and 1.2 for the FO and RO diets, respectively.

### *Growth*

At the beginning of the trial fish had a mean weight of 2053g. At the end of the trial all groups showed good performance, mean weight ranging from 3340.2g to 3664.2g, for HP-RO and MP-RO, respectively (Table 3). The oil source had a significant effect (two-way ANOVA,  $P < 0.05$ ) on growth; specifically fish fed the RO diets had higher final weight, SGR and TGC compared to fish fed the FO diets. There was no significant effect due to the protein level and no significant interactions were shown. FCR was not affected by any of the factors and varied from 0.99 to 1.10. Lastly, PPV was significantly affected by both factors. LP diets had significantly higher PPV than the other two groups (0.41, 0.43 and 0.47, for HP, MP and LP, respectively), while the RO groups had higher PPV compared to the FO groups (0.42 vs. 0.46, for FO and RO, respectively). These results are described in detail in Karalazos et al., 2010 Aquaculture, submitted for publication.

### *Tissue fatty acid compositions*

The total lipid content and the fatty acid composition of muscle and liver from fish fed the six experimental diets for 10 weeks are shown in Table 4 and 5, respectively. The muscle total lipid varied from 138.0 to 156.5 mg lipid g<sup>-1</sup> tissue. The liver lipid content was much lower, compared to that of muscle, ranging from 50.1 to 64.9 mg lipid g<sup>-1</sup> tissue. Neither the dietary protein level nor the RO inclusion affected the muscle and liver lipid contents and no significant interactions were shown by two-way ANOVA.

Regarding the FA composition of muscle and liver, they were significantly affected by the RO but not by the protein level and no significant interactions between the two factors were shown. Specifically, the inclusion of RO resulted in a significant increase of 18:1n-9, total monoenes, 18:2n-6, 20:2n-6, total n-6 PUFA and 18:3n-3. On the other hand, all saturates, including total saturated FA, 16:1n-7, 22:1, AA, EPA, DHA, total n-3 FA and the n-3/n-6 ratio were significantly reduced when the fish were fed the diets containing RO.

Notably in muscle, EPA was reduced by half (10.4 vs. 5.2%, for FO vs. RO, respectively), while the reduction in DHA was more moderate (7.8 vs. 5.0% for FO vs. RO, respectively). 18:1n-9 increased from 20.9% to 35.1%, 18:2n-6 from 6.6% to 11.6% and 18:3n-3 from 2.0% to 4.6% for FO and RO groups, respectively. Similarly in liver, EPA was reduced from 16.0% to 9.9% and DHA from 16.2% to 13.2% for FO and RO groups, respectively. The increase between FO and RO groups for 18:1n-9, 18:2n-6 and 18:3n-3 was 13.8% vs. 27.9%, 1.9% vs. 7.6% and 0.5% v. 2.9%, respectively.

The differences ( $\Delta$ ) between diet and muscle fatty acid concentrations for the six experimental diets are shown in Table 6, where negative  $\Delta$  values indicate lower values in muscle compared with diet, whereas positive values indicate accumulation in tissues relative to diet. Thus, the saturated FA, ARA, EPA, and the total n-3 PUFA were utilized to a higher extent by the fish fed the FO diets compared to the RO groups. DHA appeared to be slightly utilized in the FO groups but was accumulated in the muscle in the RO groups. On the contrary, 18:1n-9, 18:2n-6 and 18:3n-3 were found in higher concentrations in the muscle in the FO groups but were utilised in the RO groups. Likewise, the differences ( $\Delta$ ) between diet and liver fatty acid concentrations for the six experimental diets are shown in Table 7. In liver the 14:0 and 16:0 were utilized in all groups, although lower  $\Delta$  values were found in the FO groups. Similarly to the muscle, 18:1n-9, 18:2n-6 and 18:3n-3 in liver were much more utilized in the RO groups compared to the FO ones. Lastly, 18:0, ARA and DHA were accumulated in liver in all groups.

#### *Pyloric caeca phospholipid FA composition*

Pyloric caeca phospholipid FA composition (Table 8) was significantly affected mainly by the dietary oil source, although in some cases a significant main effect due to the protein level, and/or significant interactions between the two factors were also shown (two-way ANOVA,  $P < 0.05$ ). Pyloric caeca phospholipid comprised almost half as n-3 PUFA (41.6-49.9%), mainly DHA and EPA, followed by saturated FA (24.3-30.5%), largely 16:0, and monoenes (15.0-24.0%) while n-6 PUFA were less than 9% (5.3 -9.0%). DHA was the most abundant FA varying from 18.9% to 25.8% and being affected significantly both by the oil source (FO > RO) and the protein content (reduced with lower protein content). EPA was also found at high levels (16.1-22.1%) and was affected significantly by both factors, while significant interactions were also found. As shown in Figure 1 the RO groups had a significantly lower EPA content at all protein levels, whereas EPA content was significantly higher for the LP diet compared to HP but only when FO was the oil source. Regarding total saturated FA there was a significant effect due to the oil source (FO > RO) and the same pattern was shown for 16:0. Total monoenes, and mainly 18:1n-9 were significantly increased due to the dietary inclusion of RO. However, significant interactions were found for 18:1n-9 showing an effect of protein content (higher 18:1n-9 level for the LP diet vs. HP and MP) but only for the RO diets. Total n-6, mainly 18:2n-6, were increased in fish fed the RO diet, while ARA was decreased. Noticeably, all n-6 FA were at relatively low levels. Significant interactions were shown for 18:2n-6 resulting in a significantly higher content in fish fed LP diets compared to HP and MP for the RO diets.

#### *Peroxisomal $\beta$ -oxidation capacity*

The peroxisomal palmitoyl-CoA oxidation capacity in liver, red and white muscle is shown in Table 9. The peroxisomal  $\beta$ -oxidation capacity in liver ranged from 6.6 to 12.5 pmol/min/mg protein, in red muscle from 26.7 to 36.3 pmol/min/mg protein and in white muscle from 1.1 to 1.6 pmol/min/mg protein. In liver and red muscle  $\beta$ -oxidation capacity was significantly affected by the oil source ( $P = 0.035$  and  $P = 0.034$  for liver and red muscle, respectively). Specifically, RO inclusion resulted in significantly higher  $\beta$ -oxidation in both liver (7.2 vs. 9.7, for FO and RO, respectively) and red muscle (28.2 vs. 33.7, for FO and RO, respectively). However, in white muscle there was a significant interaction of the two factors (protein level and oil source) and hence, the simple main effects of the two factors were tested, as demonstrated in Figure 2. Specifically, the HP group had a significantly higher  $\beta$ -oxidation capacity than MP and LP when FO was the oil source, whereas in contrast, the  $\beta$ -oxidation capacity in HP was lower than the other two groups when RO was included in the diet. Regarding the effects of the oil source on  $\beta$ -oxidation

capacity at the three protein levels, the ranking was FO > RO for HP, RO > FO for MP while FO and RO did not significantly differ at LP.

## **Discussion**

### *Growth*

The effects and interactions of the two factors on growth are thoroughly discussed in Karalazos et. al. 2010, Aquaculture, submitted form publication. Briefly, it was shown that, in agreement with previous studies<sup>(23,30-34)</sup> no negative effects on growth and FCR were observed when the fish were fed with low protein / high lipid diets, even at protein levels below 300 g/kg. This is much lower than what had been previously tested and of significant importance regarding the tolerance in low protein diets and the utilization of lipid for energy. Moreover, the dietary inclusion of RO at expense of FO had a positive effect on final weight, SGR and TGC. Such an effect had been reported by a couple of studies<sup>(23,24)</sup> and possibly relates to the positive effect of low n-3 FA diets towards higher growth for large salmon<sup>(21,44)</sup>. This effect is probably explained by the higher digestibility of the RO, and other VO, FA, and hence better utilization of the dietary oil for energy by the fish. This theory is confirmed by the digestibility results of the present study (Karalazos et. al. 2010, Aquaculture, submitted form publication). Moreover, the increased  $\beta$ -oxidation and the changes in the pyloric caeca phospholipids discussed below could also have played a significant role.

Lastly, a positive effect of increased dietary lipid content on protein retention and, hence, on protein sparing has been previously reported<sup>(23,24,32,33)</sup> and was also confirmed in the present study as the LP diets showed a higher PPV. The inclusion of RO at the expense of FO resulted in a positive effect on PPV, also. The potential effects of dietary VO in protein sparing in fish are largely unknown, however the results of the present study are supported by the increased  $\beta$ -oxidation capacity that was shown in tissues of Atlantic salmon fed with RO diets compared to the FO diets. The results showing increased catabolism of FA for energy production may suggest a protein sparing effect. The  $\beta$ -oxidation results are discussed further below.

### *Tissue fatty acid composition*

Tissue total lipid FA composition reflects the FA composition of the diet, usually following linear correlations between the concentrations of individual FA in the diet and the tissues<sup>(16,17,19,21,24,45,46)</sup>. The results of the present study are in agreement with the previous studies showing a reduction in saturated FA, 16:1n-7, 20:4n-6, EPA, DHA and n-3/n-6 ratio, respective to the reductions in the dietary FA, with inclusion of RO. Similarly, the increase of 18:1n-9, 18:2n-6 and 18:3n-3 in the diets containing RO, compared to the FO diets, was reflected in muscle and liver

FA. It is clear that, since the FA compositions of the diets were affected by the oil source only, the changes in the muscle and liver FA compositions were also due to the dietary RO inclusion and, as expected, the dietary protein level had no significant effect on the tissue FA composition.

The changes in muscle and liver FA indicate selective utilization or retention of individual FA. It has been shown previously that when specific FA are in abundance in the diets they are selectively utilized for energy production, via  $\beta$ -oxidation, and perhaps to a lesser extent for desaturation and elongation. In contrast, when FA, and especially n-3 HUFA, are limited in the diet they are retained or deposited in the tissues<sup>(16,17,19)</sup>. For example, in the present study 18:1n-9 was increased almost 4-fold, 18:2n-6 more than 5-fold and 18:3n-3 9-fold in the diets when FO was replaced with RO, however the respective increases in muscle were less than or around 2-fold for all of the above FA, while in liver it was approx. 2-fold for 18:1n-9, less than 4-fold for 18:2n-6 and less than 6-fold for 18:3n-3, indicating selective utilization of these FA in the RO groups. On the other hand, although the reduction of EPA and DHA was more than 60% in the RO diets the decrease in muscle was approximately 50% for EPA and 35% for DHA, while in liver the reduction was approximately 35% for EPA and 20% for DHA indicating a selective retention of these FA in the tissues when the dietary supply was reduced. These results are also supported by the  $\Delta$  values for muscle and liver, that is the differences between diet and tissue fatty acid concentrations. It was shown that, when provided at high concentrations, 18:1n-9, 18:2n-6 and 18:n-3 were highly utilised in both muscle and liver and EPA and DHA were accumulated in the tissues when dietary supply was reduced. In other animals, the selective retention of essential FA in specific tissues is believed to occur by a mechanism of reacylation of sn-2-monoacylglycerols by hepatic microsomal activity of monoacylglycerol acyltransferase (MGAT), during lipolysis<sup>(47)</sup>. However, the moderate reductions of EPA and DHA shown in the present study may also be partially affected by the enhanced endogenous desaturation and elongation of dietary 18:3n-3<sup>(48-50)</sup>.

Furthermore, it is noteworthy that the results of the present trial suggest that the inclusion of RO up to 60%, at the expense of FO, in diets of Atlantic salmon at various dietary protein / lipid levels caused only moderate reductions in tissue EPA and DHA. This was also shown by Karalazos et al<sup>(24)</sup> where fish were reared at low water temperatures. Reductions in n-3 HUFA affect the quality of the final product, compromise its high nutritional value for the human consumer and should be avoided<sup>(13)</sup>. Hence, such results, leading to moderate reductions of EPA and DHA, are promising for the use of VO in commercial diets, although the present trial was conducted over a short time period, which could have masked the full extent of the FA changes that could occur over the whole production cycle of salmon.

*Pyloric caeca phospholipid FA composition*

Phospholipids are of importance in lipid digestion in fish, playing a significant role in the structure of cell membranes, of lipoproteins for the transport of lipids in the blood and lymph and also in forming intra-luminal mixed micelles along with bile salts and dietary lipids<sup>(51)</sup>. Phospholipids in fish contain mainly 16:0 and 18:1n-9 at the sn-1 position and 20:5n-3 and 22:6n-3 at the sn-2 position<sup>(26)</sup>. Hence, the intestine phospholipid FA composition and the consequent alterations occurring due to dietary FA changes may affect the uptake and digestibility of lipids and ultimately the utilization of diets, especially when high lipid diets are used. Pyloric caeca is a major site in the intestine duct for nutrient uptake and the most significant section for lipid uptake after their digestion<sup>(52)</sup>. In the present study focus was given to the interactive effects of the protein/lipid level and oil source in the FA composition of pyloric caeca.

In the present study it was shown that, regardless of the dietary treatment, n-3 PUFA was the major FA group in pyloric caeca phospholipids, consisting mainly of DHA and EPA, followed by saturated FA, largely 16:0, and monoenes (15.0-24.0%) while n-6 PUFA were less than 9% (5.3 - 9.0%), which is in accordance with previous reports<sup>(26)</sup>. The high abundance of these FA could have also been enhanced by their high recovery rate into enterocyte phospholipids<sup>(53,54)</sup>.

However, the dietary changes had significant effects on the phospholipids FA, mainly due to the oil source but also due to protein/lipid level, although to a small extent, while significant interactions were also observed. Specifically, the dietary inclusion of RO at expense of FO, resulted in significant reductions of DHA, EPA, 16:0 and 20:4n-6, and significant increases of 18:1n-9 and 18:2n-6. On the other hand, the effect of the protein content was also significant in some cases with contradicting results, including 18:0 (HP < LP) and DHA (HP > LP). Lastly, significant interactions of the two factors were also revealed by two-way ANOVA for some FA, including 18:1n-9 and 18:2n-6 with a significant increase due to the protein content (HP > LP) only for the RO groups, while EPA had a significant decrease due to the protein content (HP < LP) only for the FO groups (for these FA the effect of the oil source was significant at all protein/lipid levels). These changes reflected the changes in the dietary FA compositions, similarly to the muscle and liver total lipid FA compositions, although to a relatively smaller extent. However, altering the relative proportions of FA in intestine phospholipids, as an effect of the use of vegetable oils in the diets and more interestingly of the protein/lipid level or the interactive effects of the two factors, is most likely to affect their structure and consequently their role in the digestion and uptake of lipids and nutrients. The present study showed that lipid digestibility was improved due to RO inclusion but also (for FO only) due to low protein/high lipid diets (data presented and discussed in Karalazos et. al. 2010,

Aquaculture, submitted for publication). However, the clarification of the exact mechanisms involved requires further investigation.

#### *Peroxisomal $\beta$ -oxidation capacity*

The peroxisomal  $\beta$ -oxidation activity was measured in liver, red and white muscle. Conducting the assay on-site was not possible and hence the samples of tissues had to be frozen on dry ice and transferred to the Institute of Aquaculture in Scotland where analysis took place. Therefore, the measurement of the total  $\beta$ -oxidation activity was not possible and the results obtained represent the peroxisomal  $\beta$ -oxidation capacity. Previous studies in Atlantic salmon have shown that different tissues/organs have very different  $\beta$ -oxidation capacities as a result of their unique and different energy requirements, depending on their functions<sup>(55-60)</sup>. In agreement with that, between the three tissues assessed in the present study, red muscle had the highest  $\beta$ -oxidation activity and white muscle the lowest. However, it should be noted that the  $\beta$ -oxidation capacities of these tissues, and the consequent ranking, were expressed on a tissue protein content basis. Considering that white muscle accounts for more than 60% of the total body mass of Atlantic salmon, it becomes clear that its role in energy production for the fish is the most significant<sup>(55,56)</sup>.

It is well documented that tissue  $\beta$ -oxidation capacities are affected by various factors, including the diet and especially the dietary FA composition<sup>(50,56,57,60,61)</sup>. Specific FA, such as 16:0, 18:1n-9, 22:1n-11 and 20:1n-9 are readily catabolized, although 18:3n-3, 18:2n-6 and even EPA and DHA are also good substrates for  $\beta$ -oxidation, especially when provided at high levels<sup>(56,59,62)</sup>. Hence, dietary changes, incorporating VO, could affect the  $\beta$ -oxidation capacities of tissues. However, the results of previous studies are contradictory. Tocher et al.<sup>(50)</sup> showed that  $\beta$ -oxidation capacity was not affected either by the oil content or the oil type in diets of Atlantic salmon. On the contrary, Stubhaug et al.<sup>(56)</sup> reported that dietary RO inclusion had a positive effect on  $\beta$ -oxidation. The results of the present study showed a significant increase in liver and red muscle  $\beta$ -oxidation capacities due to RO inclusion. This could explain, at least partially, the better performance that was shown for the RO groups and the enhanced protein sparing effect. However, in white muscle an interactive effect of the protein level and the oil source was shown, suggesting a higher  $\beta$ -oxidation capacity for the FO groups than the RO ones at the HP level, whereas RO groups had higher values for the other two protein levels, although the difference was significant only for the MP diet. The higher  $\beta$ -oxidation capacity of the HP-FO group could be due to the higher content of saturated FA in that diet, although if the hypothesis is correct it remains unclear why such an effect was not reflected in the other two tissues.

## Conclusions

In conclusion, the investigation of the interactive effects of dietary protein/ lipid level and FO replacement showed that, low protein / high lipid diets can be used safely in large Atlantic salmon nutrition with regard to the growth and FCR, while the inclusion of RO at the expense of FO can enhance the growth of the fish by, increased protein sparing and  $\beta$ -oxidation. In terms of the tissue FA compositions, they were significantly affected by the RO inclusion reflecting the FA composition of the diets. However, the reduction in EPA and DHA, resulting from the dietary FA changes, was only moderate and hence, the impact on the final product quality, in terms of the nutritional value for the human consumer, was limited. Further studies on the longer term use of diets are therefore warranted.

## Acknowledgements

This work was carried out with support from BioMar AS. V. Karalazos was supported by the State Scholarships Foundation (IKY), Greece.

V. K. wrote the manuscript with assistance from all other authors especially J. G. B. and E. Å. B.; E. Å. B. was responsible for all aspects of the feeding trial, with assistance from V. K. in sample collection; V. K. was responsible for all aspects of analysis, with assistance and technical advice for  $\beta$ -oxidation analysis from J. R. D. and D. R. T. All authors read and approved the findings of the study.

None of the authors had a conflict of interest.

## References

1. Sargent JR , Tacon AG (1999) Development of farmed fish: a nutritionally necessary alternative to meat. *Proc. Nutr. Soc.* **58**, 377-383.
2. Hertrampf JW , Piedad-Pascual F (2000) *Handbook on ingredients for aquaculture feeds*. Dordrecht, Netherlands: Kluwer Academic Publishers.
3. NRC (1993) *Nutrient requirements of fish*. Washington D.C.: National Academy Press.
4. Pike IH , Barlow SM (2003) Impact of fish farming on fish stocks. *Int. Aquafeed Dir.*, 24–29.
5. Tacon AGJ (2004) Use of fish meal and fish oil in aquaculture: a global perspective. *Aquatic Resources, Culture and Development* **1**, 3-14.
6. Tidwell JH , Allan GL (2002) Fish as food: aquaculture's contribution. Ecological and economic impacts and contributions of fish farming and capture fisheries. *World Aquaculture* **33**, 44-48.
7. Delgado CL, Wada N, Rosegrant MW *et al.* (2003) *Fish to 2020: Supply and demand in changing global markets*: International Food Policy Research Institute, Washington, D.C. and WorldFish Center, Penang, Malaysia.
8. Trushenski JT, Kasper CS , Kohler CC (2006) Challenges and opportunities in finfish nutrition. *North American Journal of Aquaculture* **68**, 122-140.
9. Bell JG, McGhee F, Dick JR *et al.* (2005) Dioxin and dioxin-like polychlorinated biphenyls (PCBs) in Scottish farmed salmon (*Salmo salar*): effects of replacement of dietary marine fish oil with vegetable oils. *Aquaculture* **243**, 305-314.
10. SCAN (2000) *Dioxin contamination of feedingstuffs and their contribution to the contamination of food of animal origin. Opinion of the Scientific Committee on Animal Nutrition, adopted on November 6th 2000*. Brussels, Belgium: European Commission for Health and Consumer Protection Directorate - General.
11. SCF (2001) *Update of the Risk assessment of dioxins and dioxin-like PCBs in food based on new scientific information available since adoption of the SCF opinion of 22nd November 2000. Opinion of the Scientific Committee on Food, adopted on May 30th 2001*. European Commission for Health and Consumer Protection Directorate - General: Brussels, Belgium.
12. Bethune C, Seierstad SL, Seljeftot I *et al.* (2006) Dietary intake of differently fed salmon: a preliminary study on contaminants. *Eur. J. Clin. Invest.* **36**, 193-201.
13. Bell JG , Waagbø R (2008) Safe and Nutritious Aquaculture Produce: Benefits and Risks of Alternative Sustainable Aquafeeds. In *Aquaculture in the Ecosystem*, pp. 185-225 [M Holmer, K Black, CM Duarte, N Marbà and I Karakassis, editors]. Netherlands: Springer.
14. Turchini GM, Torstensen BE , Ng W-K (2009) Fish oil replacement in finfish nutrition. *Reviews in Aquaculture* **1**, 10-57.

15. Glencross BD (2009) Exploring the nutritional demand for essential fatty acids by aquaculture species. *Reviews in Aquaculture* **1**, 71-124.
16. Bell JG, McEvoy J, Tocher DR *et al.* (2001) Replacement of fish oil with rapeseed oil in diets of Atlantic salmon (*Salmo salar*) affects tissue lipid compositions and hepatocyte fatty acid metabolism. *J. Nutr.* **131**, 1535-1543.
17. Bell JG, McGhee F, Campbell PJ *et al.* (2003) Rapeseed oil as an alternative to marine fish oil in diets of post-smolt Atlantic salmon (*Salmo salar*): changes in flesh fatty acid composition and effectiveness of subsequent fish oil "wash out". *Aquaculture* **218**, 515-528.
18. Ng W-K, Sigholt T, Bell JG (2004) The influence of environmental temperature on the apparent nutrient and fatty acid digestibility in Atlantic salmon (*Salmo salar* L.) fed finishing diets containing different blends of fish oil, rapeseed oil and palm oil. *Aquacult. Res.* **35**, 1228-1237.
19. Torstensen BE, Frøyland L, Lie Ø (2004) Replacing dietary fish oil with increasing levels of rapeseed oil and olive oil - effects on Atlantic salmon (*Salmo salar* L.) tissue and lipoprotein lipid composition and lipogenic enzyme activities. *Aquacult. Nutr.* **10**, 175-192.
20. Torstensen BE, Frøyland L, Ørnsrud R *et al.* (2004) Tailoring of a cardioprotective muscle fatty acid composition of Atlantic salmon (*Salmo salar*) fed vegetable oils. *Food Chem.* **87**, 567-580.
21. Torstensen BE, Bell JG, Rosenlund G *et al.* (2005) Tailoring of Atlantic salmon (*Salmo salar* L.) flesh lipid composition and sensory quality by replacing fish oil with a vegetable oil blend. *J. Agric. Food Chem.* **53**, 10166-10178.
22. Caballero MJ, Obach A, Rosenlund G *et al.* (2002) Impact of different dietary lipid sources on growth, lipid digestibility, tissue fatty acid composition and histology of rainbow trout, *Oncorhynchus mykiss*. *Aquaculture* **214**, 253-271.
23. Bendiksen EÅ, Berg OK, Jobling M *et al.* (2003) Digestibility, growth and nutrient utilisation of Atlantic salmon parr (*Salmo salar* L.) in relation to temperature, feed fat content and oil source. *Aquaculture* **224**, 283-299.
24. Karalazos V, Bendiksen EA, Dick JR *et al.* (2007) Effects of dietary protein, and fat level and rapeseed oil on growth and tissue fatty acid composition and metabolism in Atlantic salmon (*Salmo salar* L.) reared at low water temperatures. *Aquacult. Nutr.* **13**, 256-265.
25. Henderson RJ, Sargent JR (1985) Chain-length specificities of mitochondrial and peroxisomal  $\beta$ -oxidation of fatty acids in livers of rainbow trout (*Salmo gairdneri*). *Comp. Biochem. Physiol. B* **82**, 79-85.
26. Sargent JR, Tocher DR, Bell JG (2002) The lipids. In *Fish Nutrition*, 3rd ed., pp. 181-257 [JE Halver and RW Hardy, editors]. San Diego, California, USA: Academic Press, Elsevier Science.
27. Kiessling KH, Kiessling A (1993) Selective utilization of fatty acids in rainbow trout (*Oncorhynchus mykiss* Walbaum) red muscle mitochondria. *Can. J. Zool.* **71**, 248-251.

- 537 28. Cho CY , Bureau DP (1997) Reduction of waste output from salmonid aquaculture through  
538 feeds and feeding. *Prog. Fish-Cult.* **59**, 155-160.
- 539 29. Halver JE , Hardy RW (2002) Nutrient flow and retention. In *Fish Nutrition*, 3rd ed., pp. 768-  
540 769 [JE Halver and RW Hardy, editors]. San Diego, California, USA: Academic Press, Elsevier  
541 Science.
- 542 30. Azevedo PA, Leeson S, Cho CY *et al.* (2004) Growth, nitrogen and energy utilization of  
543 juveniles from four salmonid species: diet, species and size effects. *Aquaculture* **234**, 393-414.
- 544 31. Azevedo PA, Leeson S, Cho CY *et al.* (2004) Growth and feed utilization of large size rainbow  
545 trout (*Oncorhynchus mykiss*) and Atlantic salmon (*Salmo salar*) reared in freshwater: diet and  
546 species effects, and responses over time. *Aquacult. Nutr.* **10**, 401-411.
- 547 32. Einen O , Roem AJ (1997) Dietary protein/energy ratios for Atlantic salmon in relation to fish  
548 size: growth, feed utilization and slaughter quality. *Aquacult. Nutr.* **3**, 115-126.
- 549 33. Hillestad M, Johnsen F, Austreng E *et al.* (1998) Long-term effects of dietary fat level and  
550 feeding rate on growth, feed utilization and carcass quality of Atlantic salmon. *Aquacult. Nutr.* **4**,  
551 89-97.
- 552 34. Solberg C (2004) Influence of dietary oil content on the growth and chemical composition of  
553 Atlantic salmon (*Salmo salar*). *Aquacult. Nutr.* **10**, 31-37.
- 554 35. Bendiksen EÅ, Arnesen AM , Jobling M (2003) Effects of dietary fatty acid profile and fat  
555 content on smolting and seawater performance in Atlantic salmon (*Salmo salar* L.). *Aquaculture*  
556 **225**, 149-163.
- 557 36. Bendiksen EÅ , Jobling M (2003) Effects of temperature and feed composition on essential fatty  
558 acid (n-3 and n-6) retention in Atlantic salmon (*Salmo salar* L.) parr. *Fish Physiol. Biochem.* **29**,  
559 133-140.
- 560 37. Folch J, Lees M , Sloane-Stanley GH (1957) A simple method for the isolation and purification  
561 of total lipides from animal tissues. *J. Biol. Chem.* **226**, 497-509.
- 562 38. Christie WW (1982) *Lipid Analyses*. 2nd ed. Oxford England: Pergamon Press.
- 563 39. Tocher DR , Harvie DG (1988) Fatty acid compositions of the major phosphoglycerides from  
564 fish neural tissues; (n-3) and (n-6) polyunsaturated fatty acids in rainbow trout (*Salmo gairdneri*)  
565 and cod (*Gadus morhua*) brains and retinas. *Fish Physiol. Biochem.* **5**, 229-239.
- 566 40. Ackman RG (1980) Fish Lipids, part 1. In *Advances in Fish Science and Technology*, pp. 86-  
567 103 [JJ Connell, editor]. Farnham, U.K.: Fishing New Books Ltd.
- 568 41. Frøyland L, Asiedu DK, Vaagenes H *et al.* (1995) Tetradecylthioacetic acid incorporated into  
569 very low density lipoprotein: changes in the fatty acid composition and reduced plasma lipids in  
570 cholesterol-fed hamsters. *J. Lipid Res.* **36**, 2529-2540.

571 42. Lowry OH, Rosebrough NJ, Farr AL *et al.* (1951) Protein measurement with the folin phenol  
572 reagent. *J. Biol. Chem.* **193**, 265-275.

573 43. Zar JH (1999) *Biostatistical analysis*. 4th ed. London, U.K.: Prentice-Hall International  
574 Editions.

575 44. Menoyo D, Lopez-Bote CJ, Bautista JM *et al.* (2003) Growth, digestibility and fatty acid  
576 utilization in large Atlantic salmon (*Salmo salar*) fed varying levels of n-3 and saturated fatty acids.  
577 *Aquaculture* **225**, 295-307.

578 45. Bell JG, Tocher DR, Henderson RJ *et al.* (2003) Altered fatty acid compositions in Atlantic  
579 salmon (*Salmo salar*) fed diets containing linseed and rapeseed oils can be partially restored by a  
580 subsequent fish oil finishing diet. *J. Nutr.* **133**, 2793-2801.

581 46. Rosenlund G, Obach A, Sandberg MG *et al.* (2001) Effect of alternative lipid sources on long-  
582 term growth performance and quality of Atlantic salmon (*Salmo salar* L.). *Aquacult. Res.* **32**  
583 (suppl. 1), 323-328.

584 47. Xia T, Mostafa N, Bhat BG *et al.* (1993) Selective retention of essential fatty acids: the role of  
585 hepatic monoacylglycerol acyltransferase. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **265**, R414-  
586 419.

587 48. Tocher DR, Bell JG, Dick JR *et al.* (2003) Effects of dietary vegetable oil on Atlantic salmon  
588 hepatocyte fatty acid desaturation and liver fatty acid compositions. *Lipids* **38**, 723-732.

589 49. Tocher DR, Bell JG, MacGlaughlin P *et al.* (2001) Hepatocyte fatty acid desaturation and  
590 polyunsaturated fatty acid composition of liver in salmonids: effects of dietary vegetable oil. *Comp.*  
591 *Biochem. Physiol. B* **130**, 257-270.

592 50. Tocher DR, Bell JG, McGhee F *et al.* (2003) Effects of dietary lipid level and vegetable oil on  
593 fatty acid metabolism in Atlantic salmon (*Salmo salar* L.) over the whole production cycle. *Fish*  
594 *Physiol. Biochem.* **29**, 193-209.

595 51. Tocher DR, Bendiksen EA, Campbell PJ *et al.* (2008) The role of phospholipids in nutrition and  
596 metabolism of teleost fish. *Aquaculture* **280**, 21-34.

597 52. Denstadli V, Vegusdal A, Krogdahl Å *et al.* (2004) Lipid absorption in different segments of the  
598 gastrointestinal tract of Atlantic salmon (*Salmo salar* L.). *Aquaculture* **240**, 385-398.

599 53. Oxley A, Tocher DR, Torstensen BE *et al.* (2005) Fatty acid utilisation and metabolism in  
600 caecal enterocytes of rainbow trout (*Oncorhynchus mykiss*) fed dietary fish or copepod oil. *Biochim.*  
601 *Biophys. Acta* **1737**, 119-129.

602 54. Pérez JA, Rodríguez C, Henderson RJ (1999) The uptake and esterification of radiolabelled  
603 fatty acids by enterocytes isolated from rainbow trout (*Oncorhynchus mykiss*). *Fish Physiol.*  
604 *Biochem.* **20**, 125-134.

605 55. Frøyland L, Lie O , Berge RK (2000) Mitochondrial and peroxisomal beta-oxidation capacities  
606 in various tissues from Atlantic salmon *Salmo salar*. *Aquacult. Nutr.* **6**, 85-89.

607 56. Stubhaug I, Frøyland L , Torstensen BE (2005)  $\beta$ -Oxidation capacity of red and white muscle  
608 and liver in Atlantic salmon (*Salmo salar* L.) - effects of increasing dietary rapeseed oil and olive  
609 oil to replace capelin oil. *Lipids* **40**, 39-47.

610 57. Stubhaug I, Lie Ø , Torstensen BE (2007) Fatty acid productive value and  $\beta$ -oxidation capacity  
611 in Atlantic salmon (*Salmo salar* L.) fed on different lipid sources along the whole growth period.  
612 *Aquacult. Nutr.* **13**, 145-155.

613 58. Tocher DR, Fonseca-Madriral J, Bell JG *et al.* (2002) Effects of diets containing linseed oil on  
614 fatty acid desaturation and oxidation in hepatocytes and intestinal enterocytes in Atlantic salmon  
615 (*Salmo salar*). *Fish Physiol. Biochem.* **26**, 157-170.

616 59. Henderson RJ , Tocher DR (1987) The lipid composition and biochemistry of freshwater fish.  
617 *Prog. Lipid Res.* **26**, 281-347.

618 60. Torstensen BE, Lie Ø , Frøyland L (2000) Lipid metabolism and tissue composition in Atlantic  
619 salmon (*Salmo salar* L.) - effects of capelin oil, palm oil, and oleic acid-enriched sunflower oil as  
620 dietary lipid sources. *Lipids* **35**, 653-664.

621 61. Torstensen BE , Stubhaug I (2004) beta-oxidation of 18:3n-3 in Atlantic salmon (*Salmo salar*  
622 L.) hepatocytes treated with different fatty acids. *Lipids* **39**, 153-160.

623 62. Stubhaug I, Tocher DR, Bell JG *et al.* (2005) Fatty acid metabolism in Atlantic salmon (*Salmo*  
624 *salar* L.) hepatocytes and influence of dietary vegetable oil. *Biochim. Biophys. Acta* **1734**, 277-288.

625

## List of Figures

Figure 1. Means of 18:1n-9 (a) and 20:5n-3 (b) (g/100g total fatty acids) of total phospholipids of pyloric caeca from Atlantic salmon fed the six experimental diets, in a two-way ANOVA, showing the effects of the two factors and their interaction.

*For each oil source, values denoted with different letters are significantly different; uppercase or lowercase letters correspond to FO or RO, respectively. Within each protein level the significant differences between FO and RO values are marked with an asterisk.*

Figure 2. Means of the peroxisomal  $\beta$ -oxidation capacity (pmol/min/mg protein) of white muscle from Atlantic salmon fed the six experimental diets, in a two-way ANOVA, showing the effects of the two factors and their interaction.

*For each oil source, values denoted with different letters are significantly different; uppercase or lowercase letters correspond to FO or RO, respectively. Within each protein level the significant differences between FO and RO values are marked with an asterisk.*

**Table 1.** Diet formulations, proximate compositions (g/kg) and energy content (kJ/g) of the six experimental diets fed to Atlantic salmon for 10 weeks.

	HP-FO	MP-FO	LP-FO	HP-RO	MP-RO	LP-RO
Component (g/kg)						
Fishmeal <sup>*</sup>	402	340	268	402	340	268
Oil seed and legume seed meals	181	190	190	181	190	190
Binder	135	130	190	135	130	190
Fish oil	304	330	351	122	132	141
Rapeseed oil <sup>†</sup>	0	0	0	182	198	211
Premixes <sup>‡</sup>	9	10	11	9	10	11
Composition (g/kg)						
Moisture	49	69	69	51	73	67
Dry Matter	951	931	931	949	927	933
Protein	353	338	291	345	328	296
Lipid	350	349	386	351	368	382
Ash	81	75	63	79	73	63
Gross Energy (kJ/g)	25.25	25.22	25.32	25.47	25.41	25.36
DP/DE <sup>§</sup>	15.4	14.0	12.3	13.7	13.0	12.3

<sup>\*</sup> South-American, Anchoveta oil

<sup>†</sup> European, non-GM, double-low quality rapeseed oil

<sup>‡</sup> Vitamin and mineral premixes prepared according to BioMar A/S commercial standards. Includes crystalline amino acids and Carophyl pink to provide 40mg/kg astaxanthin (DSM Roche, Basel, Switzerland)

<sup>§</sup> Digestible Protein/Digestible Energy

**Table 2.** Fatty acid compositions (g/100g total fatty acids) of the six experimental diets fed to Atlantic salmon for 10 weeks.

Fatty Acid	HP-FO	MP-FO	LP-FO	HP-RO	MP-RO	LP-RO
14:0	8.8	8.5	8.3	3.4	3.0	2.8
16:0	23.2	20.3	20.2	12.1	10.9	10.2
18:0	5.9	4.9	5.0	3.8	4.3	3.5
20:0	0.6	0.5	0.5	0.7	0.8	0.7
22:0	1.3	1.2	1.2	1.9	2.5	1.7
Total saturates <sup>*</sup>	40.4	35.9	35.7	22.0	21.7	19.1
16:1n-7	8.0	8.8	8.7	3.3	3.1	3.0
18:1n-9	9.5	10.9	11.3	37.7	37.7	40.0
18:1n-7	3.2	3.3	3.4	3.7	3.0	3.1
20:1n-9	1.5	1.4	1.3	1.6	1.5	1.5
22:1	1.7	1.7	1.5	1.3	1.1	0.9
24:1n-9	0.5	0.5	0.6	0.4	0.4	0.3
Total monoenes <sup>†</sup>	24.7	26.8	27.0	47.9	47.1	49.1
18:2n-6	2.1	2.4	2.7	12.5	13.0	13.8
20:2n-6	0.1	0.2	0.2	0.1	0.1	0.1
20:4n-6	1.1	1.1	1.1	0.4	0.5	0.4
22:5n-6	0.3	0.3	0.3	0.1	0.1	0.1
Total n-6 <sup>‡</sup>	4.1	4.6	4.8	13.2	14.0	14.8
18:3n-3	0.6	0.7	0.8	6.2	6.3	6.7
18:4n-3	2.1	2.3	2.2	0.8	0.8	0.7
20:4n-3	0.6	0.7	0.6	0.2	0.2	0.2
20:5n-3	17.6	19.0	18.8	5.8	6.0	5.8
22:5n-3	1.9	2.0	2.0	0.7	0.7	0.6
22:6n-3	8.0	8.0	7.9	3.2	3.2	3.0
Total n-3 <sup>§</sup>	30.8	32.7	32.5	16.9	17.3	17.1
Total PUFA	34.9	37.3	37.3	30.1	31.3	31.9
(n-3) / (n-6)	7.5	7.1	6.8	1.3	1.2	1.2

<sup>\*</sup> Includes 15:0

<sup>†</sup> Includes 16:1n-9 & 20:1n-7

<sup>‡</sup> Includes 18:3n-6, 20:3n-6 & 22:4n-6

<sup>§</sup> Includes 20:3n-3 & 22:4n-3

663 **Table 3.** Growth and performance of Atlantic salmon fed the six experimental diets for 10 weeks (Mean values (n 3) and standard deviations)

	HP-FO		MP-FO		LP-FO		HP-RO		MP-RO		LP-RO		TWO-WAY ANOVA <i>P</i>		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	protein	oil	prot x oil
Start Weight, g	2031.7	8.3	2097.0	9.6	2031.7	32.6	2065.3	28.5	2055.7	47.2	2038.3	18.6			
End Weight, g	3340.2	136.0	3491.1	134.2	3352.9	156.2	3591.8	158.7	3664.2	148.0	3405.7	92.1	0.085	0.032	0.483
FCR <sup>*</sup>	1.07	0.06	1.10	0.09	1.06	0.05	0.99	0.05	1.02	0.05	1.09	0.11	0.587	0.262	0.376
SGR <sup>†</sup>	0.86	0.07	0.88	0.07	0.86	0.08	0.95	0.05	0.99	0.03	0.88	0.06	0.262	0.025	0.422
TGC <sup>‡</sup>	3.41	0.29	3.54	0.32	3.44	0.34	3.85	0.25	4.04	0.17	3.53	0.26	0.202	0.021	0.414
PPV <sup>§</sup>	0.40	0.03	0.41	0.01	0.44	0.02	0.43	0.01	0.44	0.02	0.51	0.06	0.003	0.003	0.294

664

665 <sup>\*</sup> Feed Conversion Ratio

666 <sup>†</sup> Specific Growth Rate

667 <sup>‡</sup> Thermal Growth Coefficient

668 <sup>§</sup> Protein Productive Value

669

**Table 4.** Total lipid (mg lipid/g tissue) and fatty acid compositions (g/100g total fatty acids) of muscle from Atlantic salmon fed the experimental diets for 10 weeks (Mean values (n 3) and standard deviations)

	HP-FO		MP-FO		LP-FO		HP-RO		MP-RO		LP-RO		TWO-WAY ANOVA <i>P</i>		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	protein	oil	prot x oil
Total Lipid (mg lipid/g tissue)	142.1	14.2	142.2	24.5	138.0	9.7	147.5	13.3	156.5	14.1	143.8	3.8	0.621	0.242	0.843
<i>Fatty acid</i>															
14:0	5.8	1.3	5.6	0.4	5.7	0.3	3.4	0.1	3.2	0.1	3.4	0.3	0.766	0.000	0.988
16:0	17.4	2.9	16.3	1.3	17.0	1.5	12.4	0.4	11.9	0.2	12.3	0.7	0.677	0.000	0.941
18:0	4.0	0.7	3.7	0.4	3.8	0.4	3.1	0.1	3.1	0.0	3.1	0.2	0.695	0.001	0.903
Total saturated*	28.6	4.8	26.9	2.7	28.3	3.0	20.5	1.6	19.7	1.1	20.5	2.1	0.705	0.000	0.957
16:1n-7	6.9	0.4	7.3	0.2	7.1	0.3	4.3	0.1	4.0	0.1	4.2	0.1	0.977	0.000	0.106
18:1n-9	21.3	0.6	21.3	0.9	20.2	1.1	33.7	1.5	36.3	1.4	35.3	0.7	0.131	0.000	0.090
18:1n-7	3.6	0.1	3.8	0.3	3.8	0.3	3.4	0.4	3.0	0.2	3.4	0.1	0.427	0.004	0.214
20:1n-9	2.7	0.1	2.8	0.0	2.5	0.2	2.9	0.2	3.0	0.1	2.8	0.1	0.033	0.004	0.632
22:1	2.5	0.2	2.5	0.2	2.2	0.2	2.1	0.2	2.0	0.2	2.0	0.2	0.147	0.001	0.489
24:1n-9	0.5	0.1	0.6	0.1	0.5	0.1	0.4	0.1	0.4	0.0	0.5	0.2	0.571	0.425	0.216
Total monoenes <sup>†</sup>	37.7	0.4	38.6	0.6	36.7	1.7	47.1	1.9	49.0	1.5	48.4	1.0	0.190	0.000	0.336
18:2n-6	6.8	0.6	6.5	0.4	6.6	0.0	11.3	0.1	11.7	0.2	11.8	0.4	0.862	0.000	0.177
20:2n-6	0.4	0.1	0.4	0.0	0.4	0.0	0.6	0.0	0.7	0.0	0.6	0.1	0.194	0.000	0.484
20:3n-6	0.2	0.0	0.3	0.0	0.2	0.0	0.2	0.0	0.2	0.0	0.2	0.0	0.224	0.000	0.308
20:4n-6	0.7	0.1	0.8	0.1	0.7	0.1	0.4	0.0	0.4	0.0	0.4	0.0	0.875	0.000	0.476
Total n-6 <sup>‡</sup>	8.5	0.8	8.4	0.3	8.3	0.1	12.8	0.2	13.2	0.2	13.3	0.5	0.797	0.000	0.379
18:3n-3	2.1	0.3	2.0	0.2	2.0	0.0	4.6	0.1	4.7	0.1	4.6	0.3	0.924	0.000	0.637
18:4n-3	1.3	0.1	1.4	0.1	1.4	0.1	0.8	0.0	0.8	0.0	0.8	0.1	0.855	0.000	0.475
20:4n-3	1.0	0.1	1.0	0.1	1.0	0.1	0.7	0.0	0.7	0.0	0.7	0.0	0.746	0.000	0.419
20:5n-3	9.9	1.6	10.6	1.2	10.7	0.9	5.7	0.3	5.0	0.1	5.0	0.6	0.996	0.000	0.338

22:5n-3	3.1	0.5	3.3	0.2	3.5	0.2	2.1	0.2	1.8	0.1	1.8	0.2	0.924	0.000	0.184
22:6n-3	7.6	1.5	7.7	1.0	8.0	0.6	5.4	0.5	4.8	0.4	4.7	0.8	0.924	0.000	0.574
Total n-3 PUFA <sup>§</sup>	25.2	3.8	26.1	2.7	26.7	1.9	19.6	1.2	18.1	0.6	17.8	1.9	0.981	0.000	0.451
Total PUFA	33.7	4.4	34.5	2.9	35.0	2.0	32.4	1.4	31.3	0.7	31.1	2.4	0.996	0.039	0.680
(n-3) / (n-6)	2.9	0.3	3.1	0.2	3.2	0.2	1.5	0.1	1.4	0.0	1.3	0.1	0.761	0.000	0.049

\* Includes 15:0, 20:0, 22:0

<sup>†</sup> Includes 16:1n-9 & 20:1n-7

<sup>‡</sup> Includes 18:3n-6, 22:4n-6 & 22:5n-6

<sup>§</sup> Includes 20:3n-3 & 22:4n-3

**Table 5.** Total lipid (mg lipid g<sup>-1</sup> tissue) and fatty acid compositions (g/100g total fatty acids) of liver from Atlantic salmon fed the experimental diets for 10 weeks (Mean values (n 3) and standard deviations)

	HP-FO		MP-FO		LP-FO		HP-RO		MP-RO		LP-RO		TWO-WAY ANOVA <i>P</i>		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	protein	oil	prot x oil
Total Lipid (mg lipid/g tissue)	61.4	5.1	64.9	7.1	53.8	8.1	60.7	3.3	54.9	11.0	50.1	5.8	0.094	0.182	0.535
<i>Fatty acid</i>															
14:0	2.7	0.8	2.8	0.7	2.7	0.7	1.5	0.3	1.3	0.3	1.4	0.3	0.997	0.000	0.907
16:0	14.2	1.1	14.3	1.8	15.6	1.0	10.5	0.5	11.5	1.3	12.0	1.1	0.144	0.000	0.759
18:0	7.9	0.3	7.6	1.0	7.7	0.9	5.2	0.2	5.6	0.6	5.3	0.5	0.955	0.000	0.705
Total saturated*	25.5	1.7	25.9	2.6	26.9	2.2	18.2	0.8	19.6	2.0	20.2	1.6	0.355	0.000	0.899
16:1n-7	4.6	0.6	4.9	0.6	4.4	0.7	2.4	0.0	1.9	0.2	2.0	0.2	0.467	0.000	0.269
18:1n-9	14.8	1.0	14.5	1.4	12.0	1.4	29.8	2.6	26.9	4.5	27.1	4.0	0.038	0.000	0.106
18:1n-7	4.3	0.1	4.4	0.3	3.9	0.2	3.3	0.3	2.7	0.4	3.0	0.2	0.148	0.000	0.103
20:1n-9	2.2	0.1	2.1	0.2	1.6	0.2	3.8	0.2	3.6	0.3	3.1	0.3	0.001	0.000	0.776
22:1	0.9	0.1	0.8	0.1	0.8	0.1	0.7	0.1	0.5	0.1	0.5	0.1	0.033	0.000	0.442
24:1n-9	0.6	0.1	0.7	0.2	0.6	0.1	0.5	0.0	0.5	0.1	0.6	0.1	0.834	0.164	0.739
Total monoenes <sup>†</sup>	27.7	1.3	27.7	2.4	23.5	2.4	40.7	2.8	36.3	5.1	36.3	4.5	0.125	0.000	0.467
18:2n-6	1.9	0.2	1.9	0.2	1.9	0.3	7.6	0.3	7.2	0.7	8.2	0.7	0.298	0.000	0.355
20:2n-6	0.4	0.0	0.4	0.0	0.4	0.1	1.7	0.0	1.9	0.1	1.7	0.2	0.203	0.000	0.299
20:3n-6	0.3	0.0	0.3	0.0	0.3	0.0	0.3	0.0	0.3	0.0	0.3	0.0	0.954	0.082	0.598
20:4n-6	2.7	0.2	2.4	0.3	3.0	0.3	1.7	0.3	2.0	0.5	2.1	0.5	0.196	0.001	0.329
Total n-6 <sup>‡</sup>	5.9	0.3	5.7	0.1	6.3	0.1	11.8	0.0	11.8	0.1	12.6	0.3	0.000	0.000	0.162
18:3n-3	0.5	0.0	0.6	0.1	0.5	0.1	2.9	0.1	2.7	0.4	3.1	0.3	0.391	0.000	0.352
18:4n-3	0.4	0.0	0.4	0.1	0.4	0.1	0.2	0.0	0.1	0.0	0.1	0.0	0.614	0.000	0.651
20:4n-3	1.3	0.1	1.5	0.2	1.3	0.2	0.8	0.0	0.7	0.0	0.8	0.0	0.897	0.000	0.132
20:5n-3	15.3	1.1	15.6	0.2	17.2	0.9	9.1	0.9	10.5	1.3	10.1	1.1	0.068	0.000	0.236
22:5n-3	6.9	0.4	7.7	1.1	6.6	0.6	3.3	0.1	3.2	0.2	3.0	0.1	0.033	0.000	0.364

22:6n-3	16.3	1.1	15.0	1.2	17.2	1.5	12.3	1.3	14.2	2.2	13.0	2.4	0.738	0.003	0.199
Total n-3 PUFA <sup>§</sup>	40.9	1.6	40.8	0.3	43.3	0.2	29.3	2.1	32.3	3.2	30.9	3.2	0.120	0.000	0.114
Total PUFA	46.8	1.8	46.5	0.2	49.6	0.2	41.1	2.0	44.1	3.1	43.5	3.2	0.141	0.000	0.293
(n-3) / (n-6)	7.0	0.4	7.2	0.1	6.9	0.1	2.5	0.2	2.7	0.3	2.5	0.3	0.155	0.000	0.967

683

684 \* Includes 15:0, 20:0, 22:0

685 <sup>†</sup> Includes 16:1n-9 & 20:1n-7

686 <sup>‡</sup> Includes 18:3n-6, 22:4n-6 & 22:5n-6

687 <sup>§</sup> Includes 20:3n-3 & 22:4n-3

688

689

690 **Table 6.** Differences ( $\Delta$ )<sup>\*</sup> between diet and muscle fatty acid concentrations (g/100g total fatty acids) for the six experimental treatments

Fatty Acid	HP-FO	MP-FO	LP-FO	HP-RO	MP-RO	LP-RO
14:0	-3.0	-2.9	-2.6	0.0	0.1	0.6
16:0	-5.8	-3.9	-3.2	0.3	1.0	2.1
18:0	-1.9	-1.3	-1.2	-0.6	-1.2	-0.4
Total saturates <sup>†</sup>	-11.8	-9.0	-7.4	-1.4	-1.9	1.4
16:1n-7	-1.1	-1.5	-1.6	1.0	0.9	1.2
18:1n-9	11.7	10.4	8.9	-4.0	-1.4	-4.7
Total monoenes <sup>‡</sup>	13.0	11.7	9.7	-0.8	1.9	-0.7
18:2n-6	4.7	4.1	3.9	-1.2	-1.2	-2.0
20:4n-6	-0.4	-0.4	-0.4	0.0	-0.1	0.0
Total n-6 <sup>§</sup>	4.4	3.8	3.5	-0.4	-0.8	-1.5
18:3n-3	1.5	1.3	1.2	-1.6	-1.6	-2.1
20:5n-3	-7.7	-8.4	-8.1	-0.1	-1.0	-0.8
22:6n-3	-0.4	-0.2	0.1	2.2	1.6	1.8
Total n-3 <sup>□</sup>	-5.6	-6.6	-5.8	2.7	0.8	0.7

691

692 <sup>\*</sup> Negative  $\Delta$  values indicate lower values in muscle compared with diet, whereas positive values indicate accumulation in muscle relative to diet.

693 <sup>†</sup> Includes 15:0, 20:0, 22:0

694 <sup>‡</sup> Includes 16:1n-9, 18:1n-7, 20:1n-9, 20:1n-7, 22:1 & 24:1n-9

695 <sup>§</sup> Includes 18:3n-6, 20:2n-6, 20:3n-6, 22:4n-6 & 22:5n-6

696 <sup>□</sup> Includes 18:4n-3, 20:3n-3, 20:4n-3, 22:4n-3 & 22:5n-3

697

698 **Table 7.** Differences ( $\Delta$ )<sup>\*</sup> between diet and liver fatty acid concentrations (g/100g total fatty acids) for the six experimental treatments

699

Fatty Acid	HP-FO	MP-FO	LP-FO	HP-RO	MP-RO	LP-RO
14:0	-6.2	-5.7	-5.6	-1.9	-1.7	-1.4
16:0	-9.0	-6.0	-4.6	-1.6	0.6	1.7
18:0	2.0	2.7	2.8	1.5	1.3	1.7
Total saturates <sup>†</sup>	-14.9	-10.0	-8.8	-3.7	-2.0	1.1
16:1n-7	-4.1	-3.9	-4.3	-1.1	-1.5	-1.3
18:1n-9	5.3	3.6	0.7	-7.9	-10.8	-13.0
Total monoenes <sup>‡</sup>	3.0	0.8	-3.5	-7.2	-10.8	-12.8
18:2n-6	-0.3	-0.5	-0.7	-4.9	-5.8	-5.7
20:4n-6	1.6	1.3	1.9	1.3	1.6	1.7
Total n-6 <sup>§</sup>	1.8	1.1	1.5	-1.5	-2.2	-2.2
18:3n-3	-0.1	-0.2	-0.3	-3.3	-3.5	-3.6
20:5n-3	-2.2	-3.4	-1.7	3.3	4.5	4.3
22:6n-3	8.3	7.0	9.3	9.1	10.9	10.1
Total n-3 <sup>□</sup>	10.1	8.1	10.9	12.4	15.0	13.8

700

701 <sup>\*</sup> Negative  $\Delta$  values indicate lower values in liver compared with diet, whereas positive values indicate accumulation in liver relative to diet.

702 <sup>†</sup> Includes 15:0, 20:0, 22:0

703 <sup>‡</sup> Includes 16:1n-9, 18:1n-7, 20:1n-9, 20:1n-7, 22:1 & 24:1n-9

704 <sup>§</sup> Includes 18:3n-6, 20:2n-6, 20:3n-6, 22:4n-6 & 22:5n-6

705 <sup>□</sup> Includes 18:4n-3, 20:3n-3, 20:4n-3, 22:4n-3 & 22:5n-3

706

707 **Table 8.** Fatty acid compositions (g/100g total fatty acids) of total phospholipids of pyloric caeca from Atlantic salmon fed the experimental diets for  
708 10 weeks (Mean values (n 3) and standard deviations)

<i>Fatty acid</i>	HP-FO		MP-FO		LP-FO		HP-RO		MP-RO		LP-RO		TWO-WAY ANOVA <i>P</i>		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	protein	oil	prot x oil
14:0	2.4	0.1	2.3	0.1	2.4	0.2	1.2	0.1	1.2	0.3	1.1	0.1	0.808	0.000	0.698
16:0	20.5	0.3	21.0	0.3	20.9	0.5	16.7	0.3	17.1	0.8	16.9	0.4	0.340	0.000	0.946
18:0	6.1	0.2	6.7	0.5	6.4	0.6	5.9	0.4	6.3	0.1	6.7	0.2	0.037	0.480	0.327
Total saturated*	29.5	0.4	30.5	0.8	30.3	0.8	24.3	0.2	25.2	1.1	25.4	0.9	0.079	0.000	0.847
16:1n-7	2.9	0.1	2.9	0.3	3.0	0.3	1.3	0.0	1.3	0.2	1.2	0.1	0.902	0.000	0.548
18:1n-9	6.2	0.4	6.2	0.2	6.2	0.3	14.6	0.6	15.2	0.4	16.5	0.3	0.006	0.000	0.004
18:1n-7	3.5	0.1	3.5	0.1	3.6	0.1	2.9	0.1	2.9	0.2	3.4	0.2	0.002	0.000	0.006
20:1n-9	0.8	0.1	0.8	0.1	0.7	0.0	1.4	0.2	1.5	0.1	1.3	0.2	0.134	0.000	0.790
22:1	0.2	0.1	0.2	0.1	0.2	0.0	0.2	0.2	0.4	0.3	0.1	0.1	0.231	0.785	0.659
24:1n-9	1.1	0.1	1.0	0.1	0.9	0.1	1.0	0.1	1.0	0.0	1.0	0.1	0.394	0.682	0.375
Total monoenes <sup>†</sup>	15.2	0.5	15.0	0.7	15.0	0.6	21.9	1.0	22.6	1.2	24.0	0.7	0.162	0.000	0.076
18:2n-6	1.1	0.1	1.1	0.1	1.3	0.2	4.6	0.1	4.6	0.2	5.2	0.2	0.000	0.000	0.008
20:2n-6	0.2	0.0	0.2	0.0	0.2	0.0	0.9	0.1	0.9	0.2	0.8	0.1	0.481	0.000	0.550
20:3n-6	0.2	0.0	0.2	0.0	0.2	0.0	0.4	0.0	0.3	0.1	0.3	0.0	0.022	0.000	0.046
20:4n-6	3.0	0.2	2.8	0.1	3.0	0.2	2.2	0.1	2.1	0.1	2.0	0.1	0.487	0.000	0.321
22:5n-6	0.6	0.0	0.6	0.0	0.6	0.0	0.5	0.0	0.5	0.0	0.5	0.0	0.109	0.000	0.854
Total n-6 <sup>‡</sup>	5.4	0.2	5.3	0.1	5.5	0.0	8.8	0.2	8.5	0.4	9.0	0.2	0.049	0.000	0.612
18:3n-3	0.3	0.0	0.3	0.0	0.3	0.0	1.9	0.0	1.8	0.1	2.0	0.1	0.008	0.000	0.054
18:4n-3	0.3	0.0	0.3	0.1	0.3	0.1	0.2	0.0	0.2	0.0	0.2	0.0	0.911	0.000	0.827
20:4n-3	0.4	0.0	0.4	0.0	0.4	0.0	0.5	0.0	0.4	0.0	0.4	0.0	0.010	0.424	0.417
20:5n-3	19.3	0.6	20.7	0.2	22.1	0.2	16.1	1.1	16.1	0.3	16.7	1.0	0.004	0.000	0.041
22:5n-3	3.8	0.2	3.7	0.1	3.5	0.3	3.5	0.1	3.6	0.1	3.2	0.1	0.003	0.024	0.464
22:6n-3	25.8	0.7	23.8	0.9	22.7	0.6	22.6	0.1	21.3	1.4	18.9	0.2	0.000	0.000	0.384
Total n-3 PUFA <sup>§</sup>	49.9	0.9	49.2	0.9	49.3	0.6	45.0	1.2	43.7	1.9	41.6	1.2	0.034	0.000	0.129
Total PUFA	55.3	0.8	54.4	0.9	54.8	0.6	53.8	1.1	52.2	2.3	50.6	1.2	0.071	0.001	0.216

709		<u>(n-3) / (n-6)</u>	9.2	0.5	9.4	0.2	9.0	0.1	5.1	0.2	5.1	0.1	4.6	0.1	0.014	0.000	0.666
710	* Includes 15:0, 20:0, 22:0																
711	† Includes 16:1n-9 & 20:1n-7																
712	‡ Includes 18:3n-6 & 22:4n-6																
713	§ Includes 20:3n-3 & 22:4n-3																
714																	
715																	

716  
717  
718  
719  
  
  
720  
721

**Table 9.** Peroxisomal  $\beta$ -oxidation capacity (pmol/min/mg protein) of liver, red and white muscle from Atlantic salmon fed the experimental diets for 10 weeks (Mean values (n 3) and standard deviations)

	HP-FO		MP-FO		LP-FO		HP-RO		MP-RO		LP-RO		TWO-WAY ANOVA <i>P</i>		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	protein	oil	prot x oil
Liver	7.2	2.6	6.6	0.9	7.7	1.8	7.8	1.1	8.7	3.7	12.5	1.9	0.121	0.035	0.288
Red Muscle	30.0	2.6	28.0	2.6	26.7	4.6	36.3	8.2	31.2	4.4	33.6	4.5	0.430	0.034	0.796
White Muscle	1.6	0.1	1.2	0.1	1.3	0.0	1.1	0.1	1.6	0.0	1.4	0.2	0.908	0.719	0.000




Figure 1.




Figure 2.