



1 lesions, gill, liver, spleen and kidney were inoculated onto a variety of bacteriological agar  
2 plates and bacteriology identification and sequencing analysis was performed on significant  
3 bacterial colonies. Atypical *Aeromonas salmonicida* (aAs) *vapA* type V was the predominant  
4 bacterial species (70/215 bacteria isolates; 32.5 % of bacteria samples – 43/117 positive  
5 individual fish; 36.8 %) isolated in this survey followed by *Vibrio* species which were the  
6 most geographically prevalent bacteria. *Photobacterium indicum/profundum* was also  
7 isolated from *L. bergylta* for the first time during this study. The collection of these bacterial  
8 isolates provides useful information for disease management. Identifying the aAs isolates  
9 involved in disease in ballan wrasse could provide vital information for improving / updating  
10 existing autogenous vaccines.

11 **Key words:** atypical *Aeromonas salmonicida*, ballan wrasse, health survey, cleaner fish

12

## 13 1. INTRODUCTION

14 Ballan wrasse (*Labrus bergylta* Ascanius, 1767) and lumpsucker (*Cyclopterus lumpus*  
15 Linnaeus, 1758) are two cleaner fish species that have been intensively used by the Atlantic  
16 salmon (*Salmo salar* L.) farming industry as an alternative means to control sea lice  
17 (*Lepeophtheirus salmonis* Krøyer, 1837). The latter is an ectoparasite of the Northern  
18 hemisphere that causes major economic and welfare implications on this aquaculture  
19 industry (Treasurer 2012, Skiftesvik et al. 2013). Initially wild wrasse species (cuckoo;  
20 *Labrus mixtus* L., corkwing; *Symphodus melops* L., goldsinny; *Ctenolabrus rupestris* L. and  
21 rockcook; *Centrolabrus exoletus* L.) were used in salmon cages. However, the demand for  
22 fish and biosecurity concerns regarding the health status of wild deployed cleaner fish along  
23 with sustainable supply of wild wrasse on cage sites has led to rearing of ballan wrasse in  
24 Scotland since 2010.

1 Ballan wrasse are known to be susceptible to bacterial (*e.g.* atypical strains of *Aeromonas*  
2 *salmonicida* (aAs) and *Vibrio* spp.) (Biering et al. 2016, Gulla et al. 2016, Brooker et al.  
3 2018), parasitic (*e.g.* amoebic gill disease (AGD) (Karlsbakk et al. 2013) and viral (*e.g.*  
4 piscine myocarditis virus; PMCV) (Scholz et al. 2018) diseases. Various *Vibrio* species  
5 (*Vibrio anguillarum*, *V. ordalii* and *V. splendidus*) have also been isolated from diseased  
6 (symptomatic to vibriosis) ballan wrasse but only *V. anguillarum* originally isolated from  
7 Atlantic salmon caused high mortalities (up to 60%) in ballan wrasse under experimental  
8 conditions (Biering et al. 2016). Thus, pathogenicity of *Vibrio* species in ballan wrasse is not  
9 clear. Atypical strains of the bacterium *Aeromonas salmonicida* (As) have also been reported  
10 during mortality events of ballan wrasse in Norway (Bornø and Gulla 2016). An additional,  
11 outer membrane - the paracrystalline surface protein (A-layer protein) - plays an important  
12 role in the infection of the host as well providing protection for the bacterium by resisting  
13 host response processes (Udey & Fryer 1978, Munn et al. 1982, Kay & Trust 1991, Daly et  
14 al. 1996). The gene that encodes this protein is known as the virulence array protein A (*vapA*)  
15 and 23 A – layer (*vapA*) types of As were identified by sequencing the hypervariable region  
16 of the gene (Gulla et al. 2016, Gulla et al. 2019). Furthermore, type V and VI found to be  
17 related with cleaner fish species *L. bergylta* and *C. lumpus* in Scotland and Norway (Gulla  
18 et al. 2016, Gull et al. 2019). Cohabitation and intraperitoneal (i.p.) injection with aAs (one  
19 strain of each type V and VI used) successfully induced disease and morbidities during  
20 experimental conditions (Biering et al. 2016). Specifically, type V was found to cause the  
21 highest morbidities, suggesting that atypical strains are virulent to the species *L. bergylta*  
22 (Biering et al. 2016).

23 Information related to mortality events including causative agents / pathogens of cleaner fish  
24 such as ballan wrasse in Scotland is limited (Treasurer 2012). Bacterial disease outbreaks  
25 have been speculated to be related with aAs on commercial sites in Scotland but there are  
26 very few reports available. Prevention of disease outbreaks through vaccination is needed

1 for the species *L. bergylta* in order to improve their welfare in aquaculture and to enable  
2 their efficient performance as cleaner fish in salmon pens. Health screening and  
3 characterisation of these bacterial pathogens is essential for successful vaccine formulation.  
4 Thus, in the current study, a real-time health survey was conducted to determine the bacterial  
5 pathogens present in both farmed ballan wrasse hatcheries and Atlantic salmon cage sites  
6 (wild and farmed fish) in Scotland between February 2016 and October 2018, in order to  
7 identify the most prevalent bacterial pathogens of ballan wrasse.

8

## 9 **2. MATERIALS AND METHODS**

### 10 **2.1. Bacterial identification**

11 Healthy and / or moribund farmed and wild ballan wrasse (> 0.5 to 900 g, n= 133) were  
12 sampled from hatcheries (n= 2) and Atlantic salmon cage sites (n= 8) in Scotland between  
13 February 2016 and October 2018. Less than half of the sampled individuals (n= 43, 32.3%)  
14 had been vaccinated with an autogenous polyvalent vaccine (Ridgeway Biologicals Ltd)  
15 which included atypical furunculosis (type V and VI). From those, 42 individuals were  
16 originating from site A and had been vaccinated by two immersions (prime; ca. 0.5 g and  
17 booster vaccination; ca. 2 g) and / or injection at ca. 15 g and all the fish in the batches from  
18 which these individuals originated from had been vaccinated with the same practice. There  
19 was one more individual that had been vaccinated however, no information has been  
20 provided and whether the rest of the cleaner fish on site had been vaccinated. Furthermore,  
21 16 (12.0%) fish were unvaccinated and the rest of the individuals (n=74, 55.6%) were of  
22 unknown vaccination status (farmed or wild origin). Swab samples from skin lesions, gills,  
23 liver, and kidney were inoculated onto Marine Agar, Tryptone Soya Agar (TSA), and TSA  
24 + 5% Defibrinated Horse Blood + 1.5% NaCl, and incubated at 22°C for 24 – 72h for primary  
25 bacterial isolation. Pure colonies were then picked on the basis of morphology,

1 predominance and prevalence, streaked onto fresh plates and incubated, as described before,  
2 for purification. Passaged isolates were then tested by Gram's staining (bioMerieux) and  
3 Catalase (catalase reagents. VWR UK)/ Oxidase (oxidase strips, Oxoid UK) tests for purity  
4 confirmation and primary identification.

## 5 **2.2. Molecular analysis**

6 Bacterial DNA was extracted using genesig® Easy DNA/RNA Extraction Kit (Genesis)  
7 according to the manufacturer's instructions. Bacterial species identification was performed  
8 on the samples by targeting the subunit B protein of DNA gyrase (topoisomerase type II) –  
9 *gyrB* gene (Yamamoto et al. 2000) and V3-V4 hypervariable region of the *16S rRNA* gene  
10 (Klindworth et al. 2013) (Table 1). PCR reactions consisted of each primer at 10 µM, 1 unit  
11 of GoTaqG2 master mix (Promega), 5 µL of DNA sample and milliQ water to reach a final  
12 reaction volume of 25 µL. The following thermal cycling conditions were used in G-storm  
13 thermocycler: 1 cycle at 95°C for 5 min, 35 cycles at 95°C for 30 sec, 55°C (*gyrB*) and 44°C  
14 (16 rRNA) for 30 sec and 73°C for 1 min, followed by 1 cycle at 73°C for 7 min. The PCR  
15 product was then purified using QIAquick PCR Purification Kit (Qiagen, Germany) as  
16 described by the manufacturer and 3.5 µL of the clean-up were mixed with 2.5 µL of each  
17 of the forward and reverse primers in a separate nuclease free Eppendorf tube and 1.5 µL of  
18 nuclease free water to reach a total volume of 7.5 µL. Products sent for sequencing to GATC  
19 (Eurofins) and obtained sequences were compared to known sequences using an in silico  
20 nucleotide alignment tool 'BLAST' (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). Isolates that  
21 were recognised as presumed aAs by the naked eye; small, friable colonies, non-motile  
22 coccobacilli (prior to 16S confirmation) or by PCR testing (16S) were then confirmed to be  
23 aAs using the A-layer membrane - *vapA* primer sets (Gulla et al. 2016) to determine the *vapA*  
24 strain type (Table 1) as described by Gulla et al. (2016). The PCR product was then purified  
25 using QIAquick PCR Purification Kit (Qiagen, Germany) and samples mixed with forward  
26 and reversed primers as described above and sent for sequencing to GATC (Eurofins).

1 Sequences were analysed with Clustal Omega at EMBL-EBI (<https://www.ebi.ac.uk/>)  
2 against the published type strain sequences.

### 3 **3. RESULTS**

4 Among 327 samples (n= 133 individual fish) collected from all sites, 192 (n= 117 individual  
5 fish) had visible colonies which were identified using biochemical (Gram staining, catalase  
6 and oxidase test) and molecular (*gyrB* and *16S rRNA* sequencing, aAs *vapA* assay) analysis  
7 (Table 2).

8 Atypical As was detected in 70 (43/117 positive individual fish; 36.8%) out of 215 bacteria  
9 isolates (32.5 % of bacteria samples). Following *vapA* gene screening the majority of the  
10 aAs colonies belonged to *vapA* type V with the exception of 2 individuals that were positive  
11 to type VI from sites E and J. Atypical As were the most prevalent of the pathogenic bacteria  
12 species during this survey followed by *Vibrio* spp. and *Aliivibrio* spp.– *Vibrio ichthyoenteri*.  
13 *Vibrio splendidus*, *Vibrio tasmaniensis* *Aliivibrio logei*, *Aliivibrio salmonicida*, *Allivibrio*  
14 *finisterrensis*– (69/215; 32.1% vibrio isolates and 55/116; 47.4% positive individuals)  
15 (Figure 1 and 2). The bacteria prevalence per site is shown in Figure 1 and 2. Atypical As is  
16 most prevalent in sites A, C and E; note that site E is not presented in a pie chart as aAs (2  
17 isolates *vapA* V) were the only bacteria recovered from a single individual in a single  
18 sampling event, while *Vibrio* spp. were the most prevalent in sites B, I and J (Figure 1 and  
19 2). The majority of aAs *vapA* type V had been isolated from liver (25) and kidney (32), while  
20 the least aAs recovery was noted from fin (5), skin (4) and gill (2) samples. Also the aAs  
21 *vapA* type VI isolates (2) were from skin, liver and kidney of deployed ballan wrasse. The  
22 54.3% of the aAs (*vapA* type V and VI) isolates recovered were from vaccinated fish (21 /43  
23 individuals; 47.7%) and the majority (20 / 21) were originating from site A. Nearly half of  
24 the vaccinated individuals (20/42; 47.6 %) were positive for the bacterium (aAs).  
25 Furthermore, 8.6% of the aAs were from non vaccinated individuals (4 / 43; 9.1%) and the

1 remaining aAs isolates (37.1%) were recovered from fish with unknown vaccination status  
2 (19 / 43; 43.2%).

3 Apart from aAs another 100 (46.0 %, 82/117 individuals; 70.0%) isolates were identified  
4 and could potential be pathogenic in farmed ballan wrasse as they are known fish pathogens.  
5 These were *Aliivibrio finisterrensis*, *Aliivibrio* sp., *Aliivibrio salmonicida*, *V. anguillarum*,  
6 *Vibrio atlanticus*, *V. ichthyoenteri*, *V. lentus*, , *V. splendidus*, *V. tasmaniensis*, *T.*  
7 *ovolythicum*, *T. soleae*, *T. diecentrachi* and *Pseudomonas putida*, *Pseudomonas*  
8 *psychrophila*, *Pseudoalteromonas* sp. and *M. viscosa*. The above were recovered from gills,  
9 fins, liver, spleen and head kidney except *T. ovolythicum*, *T. soleae*, *T. diecentrachi*,  
10 *Pseudoalteromonas* sp. and *Moritella viscosa* which were isolated only in at least one of the  
11 following skin lesions, gills and / or fins.

12 No external disease signs were noted on the fish with a few exceptions. The majority of fish  
13 sampled from site A had fin rot and fish were lethargic. Internally, in some cases, the  
14 following clinical signs were observed: granulomas in the liver and/or kidney, ascites and  
15 empty gut which in some individuals was red. A suspected atypical As outbreak was active  
16 during the samplings on site A. Vaccination status of the fish did not significantly affect  
17 external or internal gross pathology for site A. Furthermore, a single wild individual from  
18 site C had a heavy skin ulcer in the flank and 3 individuals sampled at site D had pale gills,  
19 empty guts and granulomas in the organs. Co-occurrence of aAs and *Vibrio* spp. was noted  
20 for sites A, B and C in 5, 2 and 1 individual, respectively. Bacteriology analysis also showed  
21 that the individual from site C was positive for *V. splendidus* in the liver and *P. indicum* on  
22 the skin and kidney, while from three individuals (site D) *Vibrio* spp. and *Shewanella* sp.  
23 was isolated from liver and *P. indicum* from kidney.

24 Non-pathogenic bacteria also present in the samples included: *Arthrobacter* sp., *Bacillus*  
25 *simplex*, *Chryseobacterium* sp., *Colwellia* sp., *Glaciecola punicea*, *Leucothrix mucor*,  
26 *Oleispira antartica*, *Pianococcus* sp., *Planococcus* sp., *P. indicum*, *Phot. phosphoreum*,

1 *Phot. profundum*, *Photobacterium* sp., *Polaribacter irgensii*, *Polaribacter* sp.,  
2 *Pseudoalteromonas marina*, *Pseudomonas fragi*, *Psychrobacter marinicola*, *Psychrobacter*  
3 *nivimaris*, *Psychrobacter glacicola*, *Shewanella* sp, *Staphylococcus warneri*, *Vibrio*  
4 *tapetis*. *Photobacterium indicum* was also isolated from 4 locations, sites B, C, D and J with  
5 prevalence of 21.4% (3 / 14 individuals), 26.3% (5 / 19 individuals), 20.0% (1 / 5 individuals)  
6 and 22.2% (2 / 9) respectively. The sequencing data in comparison with BLAST searches  
7 gave high species similarity (97-99%) for all the above sequences.

#### 8 **4. DISCUSSION**

9 In this study a bacteriology health survey was conducted at ballan wrasse hatcheries (n= 2)  
10 and Atlantic salmon sea sites (n= 8), where wild and farmed wrasse have been deployed in  
11 Scotland, for more than 2.5 years. The majority of the sampled ballan wrasse did not have  
12 external sign of diseases with few exceptions for fish from site A, a single wild individual  
13 sampled at site C and 3 individuals sampled at site D. The predominant pathogenic bacterial  
14 species identified after bacteriology assessment and sequencing analysis (16S rRNA and  
15 *gyrB*) was aAs *vapA* type V. In corroboration with Gulla et al. (2015) aAs type V appears  
16 here to be the most predominant strain in Scotland whereas strain type VI appears to be  
17 mainly in Norway.

18 Atypical strains of As were isolated from 6 out of 10 sites that took part in this health  
19 screening survey and the bacterium was the most prevalent in 4 out of 10 sites. The results  
20 from this survey suggest that aAs was the most prevalent bacterial species at these sites  
21 between February 2016 and October 2018. It is worth noting that the aAs *vapA* type VI  
22 isolates in this survey originated from two deployed individuals in sea cages and were  
23 speculated to be related to a secondary infection following immune suppression and /or be  
24 indicative of virulence adaptation of type VI against the host. Although currently, antibiotic  
25 treatments are successfully applied for controlling disease outbreaks in hatcheries and cage



1 sites, *As* is known carry plasmids linked with antibiotic resistance. For instance *As* resistance  
2 to oxytetracycline, tetracycline and chlorafenicol has been previously reported (Adams et al.  
3 1998, L'Abée-Lund & Sørum 2002, Sørum et al. 2003). Autogenous vaccines against  
4 atypical furunculosis are also used in cleaner fish hatcheries as licenced vaccines are not  
5 available. Further characterisation of these *vapA* types through partial and / or whole  
6 sequencing (e.g. pulsed field electrophoresis; PFGE and next generation sequencing; NGS)  
7 can be helpful on identifying differences within the *aAs* strains that belong to the same type.  
8 This information can then be used to improve/update existing autogenous vaccines.

9 In addition, interestingly, 47.7% of vaccinated individuals (21 /43 individuals) were positive  
10 for the bacterium (*aAs*). Given that the majority of the positive individuals (20 / 21) had been  
11 vaccinated in the same site (site A), there is a strong suggestion that the vaccination did not  
12 appear to prevent infection by *aAs* in these fish. Protection may be influenced by the  
13 immunisation regime used as well as the isolates included in the vaccine. Both immersion  
14 and injection vaccination were being used for ballan wrasse during the time frame of this  
15 study but little is known about the efficacy of either administration routes of the vaccine.  
16 These findings support the importance for assessing immunocompetence of ballan wrasse  
17 and vaccinating the individuals at an appropriate size so that uptake and immune  
18 responsiveness to vaccine antigens is optimal. Administration of vaccines at earlier life  
19 stages of fish can lead to immunosuppression (Joosten et al. 1995, Covello et al. 2013). The  
20 majority of the individuals sampled did not show external/gross signs of disease. However,  
21 clinical signs and histopathological changes following infection by the bacterium in ballan  
22 wrasse have not yet been described, even though experimental trials have been conducted.  
23 For instance, Biering et al. (2016) showed mortalities (75 – 89 % and 51%, respectively) in  
24 juvenile ballan wrasse (50 g) infected with *aAs* either through intraperitoneal injection or  
25 cohabitation. Currently, there are not known reports of disease in farmed Atlantic salmon  
26 related with these *aAs* strains (type V or VI) and co-infection did not occur during

1 cohabitation with diseased wrasse (Gravningen et al. 1996, Treasurer 2012). Moreover,  
2 cultured Atlantic salmon are protected against typical As as routine vaccination takes place  
3 (Sommerset et al. 2005, Midtlyng 2014).

4 Bacteria belonging to the *Vibrio* and *Aliivibrio* genus (*V. ichthyenteri*, *V. splendidus*, *V.*  
5 *tasmaniensis*, *Aliivibrio salmonicida*,) known to be pathogenic to other fish species were  
6 recovered from tissue samples of ballan wrasse in this survey in 8 out of 10 sites. *V.*  
7 *splendidus*, *A. logei*, *A. wodanis* and *V. tapetis* have also been isolated from cleaner fish in  
8 Norway (Hjeltnes et al. 2018). However, *Vibrios* are universal marine bacteria and three  
9 species, *V. splendidus*, *V. ichthyenteri* and *V. pacinii* may be part of the gut flora of ballan  
10 wrasse and goldshinny wrasse (*Ctenolabrus rupestris* L.) (Birkbeck & Treasurer 2014).  
11 Thus, isolation of *V. splendidus* and *V. ichthyenteri* during the survey may have been due  
12 to accidental eruption of the gut wall, even though there is not such report. Furthermore, the  
13 presence of *Vibrio* species in the organs (liver and kidney) may have occurred at low levels  
14 that the immune system could cope with. Nonetheless these bacteria may still pose a threat  
15 as opportunistic pathogens for ballan wrasse in commercial production or during stressful  
16 events in cage sites. Similarly, ballan wrasse experienced low (10 – 20%) or no mortalities  
17 from *V. anguillarum* isolated from ballan wrasse during bath and cohabitation challenge,  
18 while i.p. injection of an Atlantic salmon strain was more virulent (50 – 60 %) (Biering et  
19 al. 2016). On the other hand lump suckers are known to be susceptible to *V. anguillarum*, *V.*  
20 *ordalii* and *V. splendidus* (Bornø & Gulla 2016). Taking the above into consideration, it is  
21 not known if ballan wrasse can act as carriers of these bacteria and infect lumpfish during  
22 cohabitation in sea pens and *vice versa*.

23 A range of non – pathogenic bacteria known to ballan wrasse were recovered during this  
24 study. From those *V. tapetis*, *T. dicentrarchi* and *P. indicum/profundum* are worth  
25 mentioning. *Vibrio tapetis* is a known pathogen for bivalves, clam species and Atlantic  
26 halibut (Reid et al. 2003, Paillard, 2004). Although, juvenile ballan wrasse (approx. 30 g)

1 were not susceptible to these bacteria species during cohabitation challenge and only i.p.  
2 injected shedder fish experienced mortalities (Gulla et al. 2017), it is not known if larvae or  
3 younger age juvenile ballan wrasse (<30 g) can be susceptible to the bacteria under rearing  
4 conditions. *Tenacibaculum dicentrachi* was isolated from ballan wrasse during this survey  
5 and to the best of the author's knowledge this is the first time that *T. dicentrachi* was  
6 recovered from ballan wrasse in Scotland. The bacterium belongs to the Family  
7 *Flavobacteriaceae* and *Tenacibaculum* spp. are ubiquitous bacteria of the marine  
8 environment with a few members of the genus related with fish diseases. For instance, *T.*  
9 *dicentrachi* was first isolated from European sea bass (*Dicentrarchus labrax*) in Spain  
10 (Piñeiro-Vidal et al. 2012) and is now a rapidly emerging pathogen of farmed Atlantic  
11 salmon in Chile (Avendaño-Herrera et al. 2016). In Norway, isolates of the genus have been  
12 recovered from skin ulcers from salmonids and non-salmonid species (Olsen 2017).  
13 Understanding the pathogenicity of this bacterium in individual ballan wrasse is important  
14 considering that the closely related species of the genus are an emerging bacteria pathogen  
15 for salmonids. Cohabitation with diseased salmon can lead to disease transmission between  
16 hosts.

17 *Photobacterium indicum/profundum*, also reported in this screening, has not previously been  
18 associated with fish disease outbreaks but has been isolated from moribund lobster and  
19 associated with stress (Basti et al. 2011). A number of isolates (7/151; 6/82 individuals) were  
20 recovered from diseased ballan wrasse in this study which might be indicative of a secondary  
21 infection after individuals had been infected with aAs. Recently, *Photobacterium* sp., were  
22 recovered from lumpsuckers experiencing mortalities due to *Pseudomonas anguilliseptica*  
23 under rearing conditions in Scotland (Treasurer & Birkbeck 2018). Further investigation is  
24 needed regarding the pathogenicity and transmission between hosts in order to understand  
25 the importance of this bacterium in cleaner fish hatcheries and deployment sites.

1 Overall, aAs was the most prevalent bacterial species isolated from ballan wrasse on the farm  
2 sites considering the number of individuals sampled in total, followed by *Vibrio* species  
3 which were the most geographically prevalent bacteria. Understanding the prevalence of  
4 these pathogens is vital for mitigating disease outbreaks by optimising fish husbandry and  
5 biosecurity practices. Furthermore, the collection of these bacterial isolates provides useful  
6 information for disease management. Also, characterisation of the aAs *vapA* types could  
7 provide important information for improving/updating existing autogenous vaccines.

8

## 9 **Acknowledgements**

10 The authors would like to thank Ridgeway Biologicals for samples collection and culture  
11 identification and MOWI Scotland and Scottish Sea Farms for their cooperation. This study  
12 was funded by the Scottish Aquaculture Innovation Centre (SAIC) under project number  
13 SL-2015-01 'Sustainable production of wrasse for sea lice control' (Health WP3).

14

## 15 **LITERATURE CITED**

- 16 Adams CA, Austin B, Meaden PG and McIntosh D (1998) Molecular characterization of  
17 plasmid-mediated oxytetracycline resistance in *Aeromonas salmonicida*. *Appl Environ*  
18 *Microb* 64:4194–4201.
- 19 Austin B and Austin DA (2017). Bacterial Fish Pathology. In C. Aeromonadaceae  
20 Representative (*Aeromonas salmonicida*), 6th ed, Springer, pp. 216 - 217
- 21 Avendaño-Herrera R, Irgang R, Sandoval C, Moreno-Lira P, Houel A, Duchaud E, M  
22 Poblete-Morales P, Nicolas P and Ilardi P (2016) Isolation, characterization and  
23 virulence potential of *Tenacibaculum dicentrarchi* in salmonid cultures in Chile.  
24 *Transbound Emer Dis*, 63: 121–126.

- 1 Basti D, Bouchard D and Lichtenwalner A (2011) Safety of Florfenicol in the Adult Lobster  
2 (*Homarus americanus*). J Zoo Wildlife Med 42: 131-133.
- 3 Biering E, Krossøy VB, Gulla S and Colquhoun DJ (2016) Challenge models for atypical  
4 *Aeromonas salmonicida* and *Vibrio anguillarum* in farmed ballan wrasse (*Labrus*  
5 *bergylta*) and preliminary testing of a trial vaccine against atypical *Aeromonas*  
6 *salmonicida*. J Fish Dis 39:1257–1261.
- 7 Birkbeck TH and Treasurer JW (2014) *Vibrio splendidus*, *Vibrio ichthyenteri* and *Vibrio*  
8 *pacinii* isolated from the digestive tract microflora of larval ballan wrasse, *Labrus*  
9 *bergylta* Ascanius, and goldsinny wrasse, *Ctenolabrus rupestris* (L.). J Fish Dis 37:  
10 69–74.
- 11 Brooker AJ, Papadopoulou A, Gutierrez C, Rey S, Davie A and Migaud H (2018)  
12 Sustainable production and use of cleaner fish for the biological control of sea lice:  
13 recent advances and current challenges. Vet Rec 183: 383.
- 14 Bornø G and Gulla S (2016) The Health Situation in Norwegian Aquaculture: The  
15 Norwegian Veterinary Institute, p. 117.
- 16 Daly JG, Kew AK, Moore AR and Olivier G (1996) The cell surface of *Aeromonas*  
17 *almonicida* determines in vitro survival in cultured brook trout (*Salvelinus fontinalis*)  
18 peritoneal macrophages. Microb Pathog 21:447– 461.
- 19 Gravningen K., Kvenseth PG and Hovlid RO (1996) Virulence of *Vibrio anguillarum*  
20 serotypes 01 and 02, *Aeromonas salmonicida* subsp. *salmonicida* and atypical  
21 *Aeromonas salmonicida* to goldsinny wrasse. In: Wrasse Biology and use in  
22 Aquaculture (ed. by MDJ Sayer, JW Treasurer and MJ Costello), Blackwell, Oxford,  
23 pp. 247–250.

- 1 Gulla S, Lund V, Kristoffersen AB, Sørum H and Colquhoun DJ (2016) *vapA* (A-layer)  
2 typing differentiates *Aeromonas salmonicida* subspecies and identifies a number of  
3 previously undescribed subtypes. J Fish Dis 39:329–42.
- 4 Gulla S, Rønneseth A, Sørum H, Vågnes Ø, Balboa S, Romalde JL and Colquhoun DJ (2017)  
5 *Vibrio tapetis* from wrasse used for ectoparasite bio-control in salmon farming:  
6 phylogenetic analysis and serotyping. Dis Aquat Organ 125:189-197.
- 7 Gulla S, Bayliss S, Björnsdóttir B, Dalsgaard I, Haenen O, Jansson E, McCarthy U, Scholz  
8 F, Vercauteren M, Verner-Jeffreys D, Welch T, Wiklund T and Colquhoun DJ (2019)  
9 Biogeography of the fish pathogen *Aeromonas salmonicida* inferred by *vapA*  
10 genotyping. FEMS Microbiol Lett, 366.
- 11 Hjeltnes B, Bang-Jensen B, Bornø G, Haukaas A and Walde C S (Ed.) (2018) The Health  
12 Situation in Norwegian Aquaculture 2017 Norwegian Veterinary Institute..
- 13 Joosten PHM, Aviles-Trigueros M, Sorgeloos P, Rombout JHWM (1995) Oral vaccination  
14 of juvenile carp (*Cyprinus carpio*) and gilthead seabream (*Sparus aurata*) with  
15 bioencapsulated *Vibrio anguillarum* bacterin. Fish Shellfish Immunol 5:289–299
- 16 Kay WW, Buckley JT, Ishiguro EE, Phipps BM, Monette JP and Trust TJ (1981) Purification  
17 and disposition of a surface protein associated with virulence of *Aeromonas*  
18 *salmonicida*. J Bacteriol 147:1077–1084.
- 19 Kay WW and Trust TJ (1991). Form and functions of the regular surface array (S-layer) of  
20 *Aeromonas salmonicida*. Experientia 47:412–414.
- 21 Karlsbakk E, Olsen AB, Einen ACB, Mo TA, Fiksdal IU, Aase H, Kalgraff C, Skår SÅ and  
22 Hansen H (2013) Amoebic gill disease due to *Paramoeba perurans* in ballan wrasse  
23 (*Labrus bergylta*). Aquaculture 412-413: 41 – 44.

- 1 Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, and Glöckner FO (2012)  
2 Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-  
3 generation sequencing-based diversity studies. *Nucleic Acids Res* 41.
- 4 L'Abée-Lund TM and Sørum H (2002) A global non-conjugative tet C plasmid, pRAS3,  
5 from *Aeromonas salmonicida*. *Plasmid* 47, 172–181.
- 6 Lund V, Espelid S and Mikkelsen H (2003b) Vaccine efficacy in spotted wolffish  
7 *Anarhichas minor*: relationship to molecular variation in A-layer protein of atypical  
8 *Aeromonas salmonicida*. *Dis Aquatic Organ* 56, 31– 42.
- 9 Midtlyng PJ (2014) Vaccination against Furunculosis. *In Fish Vaccination*, 1<sup>st</sup> Edition, Eds.  
10 R. Gudding, A Lillehaug, Ø. Evensen, Wiley Blackwell, Ch. 16, pp. 185 – 199.
- 11 Munn CB, Ishiguro EE, Kay WW and Trust TJ (1982) Role of surface components in serum  
12 resistance of virulent *Aeromonas salmonicida*. *Infection and Immunity* 36:1069–  
13 1075.
- 14 Olsen AB, Gulla S, Steinum T, Colquhoun DJ, Nilsen H. and Duchaud E (2017) Multilocus  
15 sequence analysis reveals extensive genetic variety within *Tenacibaculum spp.*  
16 Associated with ulcers in sea-farmed fish in Norway. *Vet Microbiol* 205: 39–45.
- 17 Paillard C (2004) A short review of brown ring disease, a vibriosis affecting clams,  
18 *Ruditapes philippinarum* and *Ruditapes decussatus*. *Aquat Living Resour* 17:467–  
19 475.
- 20 Piñeiro-Vidal M, Gijón D, Zarza C and Santos Y (2012) *Tenacibaculum dicentrarchi* sp.  
21 nov., a marine bacterium of the family *Flavobacteriaceae* isolated from European  
22 sea bass. *Int J Syst Evol Micr* 62, 425-429.

- 1 Reid H, Duncan HI, Laidler A, Hunter D and Birkbeck TH (2003) Isolation of *Vibrio tapetis*  
2 from cultivated Atlantic halibut (*Hippoglossus hippoglossus* L.). *Aquaculture* 221:  
3 65–74
- 4 Scholz F, Ruane NM, Morrissey T, Marcos-López M, Mitchell S, O'Connor I, Mirimin L,  
5 MacCarthy E and Rodger HD (2018) Piscine myocarditis virus detected in corkwing  
6 wrasse (*Symphodus melops*) and ballan wrasse (*Labrus bergylta*). *J Fish Dis* 41:147  
7 – 152.
- 8 Skiftesvik, AB, Bjelland RM, Durif CMF, Johansen IS and Browman HI (2013) Delousing  
9 of Atlantic salmon (*Salmo salar*) by cultured vs. wild ballan wrasse (*Labrus*  
10 *bergylta*). *Aquaculture* 402-403:113–118.
- 11 Sommerset I, Krossøy B, Biering E and Frost P (2005) Vaccines for fish in aquaculture.  
12 *Expert Rev Vaccines* 4:89–101.
- 13 Sørum H, L'Abée-Lund TM, Solberg A and Wold A (2003) Integron-containing IncU R  
14 plasmids pRAS1 and pAr-32 from the fish pathogen *Aeromonas salmonicida*.  
15 *Antimicrob Agents Ch* 47:1285–1290.
- 16 Treasurer JW (2012) Diseases of north European wrasse (*Labridae*) and possible interactions  
17 with cohabited farmed salmon, *Salmo salar* L. *J Fish Dis* 3:555–562.
- 18 Treasurer JW and Birkbeck H (2018) *Pseudomonas anguilliseptica* associated with  
19 mortalities in lumpfish reared in Scotland. *B Eur Assoc Fish Pat* 38:2018.
- 20 Udey JL and Fryer LR (1978) Immunization of fish with bacterins of *Aeromonas*  
21 *salmonicida*. *Mar Fish Rev* 40:12–17.
- 22 Yamamoto S, Kasai H, Arnold DL, Jackson RW, Vivian A and Harayama S (2000)  
23 Phylogeny of the genus *Pseudomonas*: intrageneric structure reconstructed from the  
24 nucleotide sequences of *gyrB* and *rpoD* genes. *Microbiology* 146: 2385 – 2394.






1 Table 1 List of primers used for bacteria species identification.

Primer	Primer name	Target gene	Annealing (°C)	Application	Reference
CAGGAAACAGCTATGACCAAYGSNGG	UP -1E	<i>gyrB</i>	60	PCR	Yamamoto et al., 2000
NGGNAARTTYRA					
TGTAAAACGACGGCCAGTGCNNGRT	APrU	16S	44	PCR	Klintworth et al., 2013
CYTTYTCYTGRCA					
AGAGTTTGATCMTGGC	Bact-0008	16S	44	PCR	Klintworth et al., 2013
CCGTCAATTCMTTGGAGTTT	Bact-0907				
CTGGACTTCTCCACTGCTCA	F2	<i>vapA</i>	53	PCR and sequencing	Lund et al., 2003b
ACGTTGGTAATCGCGAAATC	R3				Gulla et al., 2016

1 Table 2. Standard bacteriology tests (Gram stain, shape, catalase, oxidase) on pathogenic  
 2 bacteria isolated from skin lesions, gills, liver and kidney swabs of moribund or recovered  
 3 ballan wrasse (*Labrus bergylta*; >0.5 to 900 g) during disease outbreaks in hatcheries and  
 4 salmon sea cage sites in Scotland between February 2016 and October 2018. Sequencing  
 5 similarity represents blasts results from *16S* and *gyrB* sequencing.

<b>Bacteria species</b>	<b>Shape</b>	<b>Catalase (-/+)</b>	<b>Oxidase (-/+)</b>	<b>Sequencing similarity (%)</b>
atypical <i>Aeromonas salmonicida</i>	Bipolar rods	-	+	99-100
<i>Vibrio spp.</i>	Curved rods	+	+	99-100
<i>Vibrio (Allivibrio) salmonicida</i>	Curved rods	+	+	99-100
<i>Vibrio tasmaniensis</i>	Rods	+	+	99
<i>Vibrio splendidus</i>	Short rods	+	+	99-100
<i>Vibrio logei</i>	Cigar like rods	+	-	99
<i>Vibrio splendidus</i>	Rods	+	+	99-100
<i>Vibrio ichthyoenteri</i>	Thin rods	+	+	96-100
<i>Vibrio sp</i>	Rods	+	+	100
<i>Vibrio anguillarum</i>	Bipolar rods	-	+	99
<i>Tenacibaculum dicentrachi</i>	Curved rods	+	+	99
<i>Tenacibaculum solea</i>	Slender rods	-	+	100
<i>Tenacibaculum ovoliticum</i>	Filamentous rods	+	-	100
<i>Pseudomonas spp.</i>	Short bipolar rods	+	+	99-100
<i>Pseudoalteromonas spp.</i>	Bipolar rods	-	+	100
<i>Moritella viscosa</i>	Slightly curved rods	+	+	99
<i>Flavobacterium frigidarium</i>	Chaining cocci-bacillus	+	-	97-100


6



1

2 **Figure 1.** Bacteria percentage recovery from 4 out of 10 sites during a health screening survey for ballan wrasse in Scotland between 2016 and 2018  
 3 (Part 1). Swabs from skin lesions, gills, liver and kidney plated on variety of agar plates (Marine Agar, Tryptone Soya Agar (TSA), and TSA + 5%

- 1 Defibrinated Horse Blood + 1.5% NaCl) for phenotypic and sequencings identification. Pie chart for Site E is not shown as only atypical *Aeromonas*
- 2 *salmonicida* was isolated in a single sampling (2/2). Site F, single sampling point with no bacteria recovery from individuals sampled.



- 3
- 4 Figure 2. Bacteria percentage recovery from 4 out of 10 sites during a health screening survey for ballan wrasse in Scotland between 2016 and 2018 (Part
- 5 2). Swabs from skin lesions, gills, liver and kidney plated on variety of agar plates (Marine Agar, Tryptone Soya Agar (TSA), and TSA + 5% Defibrinated

- 1 Horse Blood + 1.5% NaCl) for phenotypic and sequencings identification. Pie chart for Site E is not shown as only atypical *Aeromonas salmonicida* was
- 2 isolated in a single sampling (2/2). Site F, single sampling point with no bacteria recovery from individuals sampled.