



Investigating the Effect of Restaurant Menu Labelling on Consumer Food Choices Using a Field Experiment

Journal:	<i>British Food Journal</i>
Manuscript ID	BFJ-04-2021-0432.R1
Manuscript Type:	Research Paper
Keywords:	Consumer behavior, consumer choice, Menu labelling, Consumer perceptions, Calorie labelling, Traffic light coding

SCHOLARONE™
Manuscripts

Published in *British Food Journal* by Emerald. The original publication is available at: <https://doi.org/10.1108/BFJ-04-2021-0432>. This author accepted manuscript is deposited under a Creative Commons Attribution Non-commercial 4.0 International (CC BY-NC) licence. This means that anyone may distribute, adapt, and build upon the work for non-commercial purposes, subject to full attribution. If you wish to use this manuscript for commercial purposes, please contact permissions@emerald.com

Investigating the Effect of Restaurant Menu Labelling on Consumer Food Choices Using a Field Experiment

1. Introduction

Obesity and overeating are causing a global public health crisis (Pomeranz, 2011; Arno and Thomas, 2016; Crockett *et al.*, 2018; Fitzgerald *et al.*, 2018). This phenomenon is costly to healthcare systems worldwide, with the prevalence of obesity and the complications that result from poor dietary choices expected to rise (Wootan and Osborn, 2006; Wootan, Osborn and Malloy, 2006). Many developed countries have seen an increase in out-of-home eating, which has been linked to an increase in obesity and diet-related morbidities (S. N. Bleich *et al.*, 2017; Bleich, Moran and Jarlenski, 2018; Kerins *et al.*, 2018). There is a heated debate over whether governments or the food industry should spearhead new strategies to combat these issues (Almanza, Nelson, and Chai, 1997; Alexander, O'Gorman, and Woods, 2010; Fitzgerald *et al.*, 2018). Many high-income countries have voluntary government-led initiatives with varying levels of private-sector participation (Bleich, Moran and Jarlenski, 2018). Currently, in the UK and the EU, packaged food labelling is mandatory, but restaurant menu labelling is not. This is a common situation in countries that have evaluated the use of menu labelling legislation (Almanza, Nelson and Chai, 1997; Vandevijvere and Vanderlee, 2019).

A substantial amount of research has been conducted to determine what type of labelling most effectively promotes positive behaviour change toward healthy food choice and consumption. This is especially pertinent given the current political debate in the United Kingdom, Europe, and North America over whether the policy is required to address nutritional labelling on restaurant menus (Britt *et al.*, 2011; Mah *et al.*, 2013; Geaney *et al.*, 2015; Arno and Thomas, 2016). Menu labelling has gained public and legislative support (Mah *et al.*, 2013; Sinclair, Cooper, and Mansfield, 2014; Reale, Flint, and Capehorn, 2015), with many workplaces and healthcare organisations developing and implementing menu labelling policies on their own at both the national and local authority levels (Kerins *et al.*, 2018). One UK policy initiative is the public consultation launched by the Scottish Government to understand the barriers and motivators for restaurant menu labelling in Scotland (Food Standards Scotland, 2018).

Several studies have found widespread support for calorie labelling on menus (Kuo *et al.*, 2009; Mackison, Wrieden and Anderson, 2009; Kerins *et al.*, 2018). While there is public support for calorie labelling, policy issues include how to implement this labelling. In Scotland and Northern Ireland, this is accomplished by using MenuCal, an online tool that assists businesses in calculating the calorie content of menu items. Food Standards Scotland (FSS) is currently offering MenuCal as a tool for businesses to label their menus without incurring high costs (Geaney *et al.*, 2015; Fitzgerald *et al.*, 2018; Kerins *et al.*, 2018). Its goal is to encourage healthier food choices by informing people about the nutritional details of foods, specifically the calorie content and allergen content. This is especially important given the higher proportion of people who eat out of the house on a regular basis, as well as the fact that meals eaten out of the house are often higher in calories and certain nutrients, such as salt

1
2
3 and fats, than those cooked at home (Orfanos *et al.*, 2009; Kerins *et al.*, 2018). While research
4 on menu labelling shows that calorie labelling has benefits, there is no agreed-upon form of
5 labelling and its potential impact on consumer choices.
6

7
8 This research adds to the literature by investigating the effect of menu labelling on
9 consumers' actual meal purchases in a real-world setting using a field experiment at a
10 restaurant. In addition, we are doing something different with the menu labels. We use a
11 combination of traffic-light colour coding and a pictogram to help consumers understand the
12 levels of nutrients and calorie content, respectively. Although several studies have looked into
13 menu labelling on options, with the main focus being primarily on calorie labelling, how to
14 communicate calories and other nutrient information on menu labels has been overlooked,
15 particularly because real-world research in this area is limited. Thus, in this study, we
16 contribute to a better understanding of the impact of menu labelling on actual meal
17 purchases, as well as the best way to communicate calorie and nutrient information to
18 consumers, and we share our experience designing a field experiment with a restaurateur for
19 future research.
20
21
22
23

24 The rest of the paper is organised as follows. Section 2 provides a brief review of menu
25 labelling. Section 3 presents the study design, data and analysis. Section 4 reports the main
26 results of the field experiment. Section 5 discusses the findings and limitations.
27 Finally, Section 6 concludes and suggests avenues for further research.
28
29

30 **2. Menu Labelling**

31 **2.1. Benefits of Menu Labelling**

32
33 The proponents of menu labelling highlight two potential sources of improvement in public
34 health as a result of menu labelling. The first is that people will make healthier choices and
35 consume fewer calories on average (Bassett *et al.*, 2008; S. N. Bleich *et al.*, 2017; Crockett *et*
36 *al.*, 2018). While menu labelling has been shown to nudge consumers to make healthier
37 choices (Arno and Thomas, 2016; Cawley, Susskind, and Willage, 2020), not all studies have
38 produced statistically significant results to support this claim (Sinclair, Cooper, and Mansfield,
39 2014). There is frequently a small decrease in the mean number of calories ordered before
40 and after menu labels are introduced. However, this could still result in significant reductions
41 in healthcare spending (Bassett *et al.*, 2008; Thomas, 2015). There is growing evidence in the
42 Republic of Ireland that introducing a menu labelling initiative can influence consumer choice
43 to promote healthier options (Food Safety Authority of Ireland, 2012; Kelly *et al.*, 2014). Many
44 studies have shown that a sizable percentage of people would use menu labelling at least
45 some of the time (Auchincloss *et al.*, 2013; Hammond *et al.*, 2013; Geaney *et al.*, 2015).
46
47
48
49
50
51

52 Another effect of calorie menu labelling would be that restaurants would redesign their
53 menus to include healthier options (Auchincloss *et al.*, 2014; Condrasky *et al.*, 2015; Crockett
54 *et al.*, 2018; Vandevijvere and Vanderlee, 2019; Cawley, Susskind and Willage, 2020). There is
55 a lot of evidence to support the idea that businesses self-regulate and reformulate menu
56 items to reduce calories in products after menu labelling is implemented (Bruemmer *et al.*,
57 2012; Bleich, Moran and Jarlenski, 2018). Across studies on restaurateurs' behaviour, there is
58 a consistent trend indicating that they offer products with fewer calories on menus (Harnack
59
60

1
2
3 and French, 2008; Restrepo, 2017). Large chain restaurants appeared to be part of this trend,
4 although 'core' items did not change calorie count over time, but newly introduced items
5 appeared to have lower calories than the items they replaced (Bleich, Wolfson and Jarlenski,
6 2016). This caveat was discovered when studies found no changes in median calorie content
7 while also seeing a decrease in average calorie content, indicating that there may still be room
8 to improve the healthy options available (Bleich, Moran and Jarlenski, 2018).
9

10
11 Some evidence suggests that businesses are not averse to reshaping menu items (Obbagy *et*
12 *al.*, 2011). In England, the Responsibility Deal emphasised that taking voluntary responsibility
13 to reformulate menu items could be part of a strategy to reduce high-calorie consumption
14 outside the home (Nikolaou, Lean and Hankey, 2014). While it was acknowledged that
15 reformulation alone is unlikely to have a significant impact on public health, it is critical to any
16 public health diet strategy (Crockett *et al.*, 2018; Kerins *et al.*, 2018). According to studies
17 from the United States, where menu labelling is required, many restaurants reduced their
18 mean calorie count due to reformulation (Bruemmer *et al.*, 2012; S. N. Bleich *et al.*, 2017).
19 While there is mixed evidence from the Responsibility Deal in England regarding voluntarily
20 providing menu labels, growing evidence suggests that a voluntary policy measure may not
21 be sufficient (Geaney *et al.*, 2015; Knai *et al.*, 2015).
22
23

24 **2.2. Barriers to Menu labelling**

25
26 While labelling policies exist in the food processing and manufacturing industries, the
27 restaurant industry faces unique operational challenges when it comes to implementing
28 menu labels. The cost to the industry is one such barrier. According to the British Hospitality
29 Association, menu labelling would benefit public health little while posing high costs to
30 businesses, which could be passed on to consumers (Thomas, 2015). Another common barrier
31 mentioned in the literature is that restaurant owners and employees do not believe they have
32 the necessary skills or knowledge to accurately measure the calories of menu items (Obbagy
33 *et al.*, 2011; Kerins *et al.*, 2018). This contributes to the cost of implementing calorie labelling
34 on restaurant menus, as outside companies are frequently hired to provide this service
35 (Geaney *et al.*, 2015; Kerins *et al.*, 2018). Without the expertise required to accurately
36 calculate calorie counts, menu item innovation and creation may be stifled. This is especially
37 problematic for small or independent restaurants that cannot afford to pay for accurate
38 calorie information for new menu items on a regular basis (Condrasky *et al.*, 2015; Thomas,
39 2015). Other practical issues for menu labelling include a lack of menu space and thus
40 difficulty in designing aesthetically pleasing menus, a lack of standardised cooking methods
41 in some restaurants, which could lead to inconsistent portion sizes and, as a result, misleading
42 menu label information, and offering no menus or portion control at buffet-style dining
43 (Alexander, O'Gorman, and Woods, 2010; Auchincloss *et al.*, 2013; Yepes, 2015).
44
45
46
47
48
49
50
51
52

53 The desire to avoid information overload for customers on the menu, as well as the fear of
54 accidentally providing misleading information on menus, have been cited as reasons why many
55 restaurants in the UK have not participated in menu labelling initiatives (Knai *et al.*, 2015;
56 Fitzgerald *et al.*, 2018). According to research, more information about the benefits of menu
57 labelling should be provided to restaurant owners, as well as practical assistance in nutritional
58
59
60

1
2
3 analysis, nutrition training for staff, and menu design from public health practitioners and
4 academics (Almanza, Nelson and Chai, 1997; Fitzgerald *et al.*, 2018).
5

6 **2.3. Menu Labels Used**

7
8 One aspect of the debate over nutritional labelling on restaurant menus that is frequently
9 overlooked is what format of menu labelling would be most beneficial. While the Scottish
10 Government considers whether to require calorie labelling on menus, some research suggests
11 that different variations of menu labelling can have varying levels of usefulness for
12 consumers. A recent Cochrane review (Crockett, 2018) highlighted a variety of menu labelling
13 interventions, ranging from simply requiring the labelling of calorie content (Bollinger, Leslie,
14 and Sorensen, 2011) and requiring a statement of recommended calorie intake (Girz *et al.*,
15 2012) to the labelling of other nutritional information such as fat, salt, and sugar content, as
16 well as the calorie content of each menu (Hammond *et al.*, 2013). There are very few
17 examples of mandatory nutritional labels other than calorie labelling. Menu boards in New
18 York, USA, for example, are required to display the sodium content of each item (Vandevijvere
19 and Vanderlee, 2019). Various menu labelling interventions have also been used in studies to
20 compare a variety of possible outcomes, such as calories provided as numbers and calories
21 represented with traffic-light codes for low, medium, and high contents (Hammond *et al.*,
22 2013). However, the results between different menu-labelling practices can be inconclusive
23 (Ellison, Lusk, and Davis, 2013; Liu *et al.*, 2012), with some evidence of symbolic labels being
24 beneficial (Holmes *et al.*, 2013; Antonelli and Viera, 2015).
25
26
27
28
29
30

31 In our study, we design and implement a new menu label that communicates calorie content
32 and nutrients (fat, saturated fat, sugars, and salt) using traffic-light colour coding and a
33 pictogram. The traffic light colour coding helps explain nutrient levels (low, medium, and
34 high), while the pictogram aids understanding of calorie information by providing walking
35 minutes to burn associated calories. Previous research has found that physical-activity
36 equivalent calorie labelling may influence food choices by promoting healthy options or
37 decreasing intentions to consume unhealthy options (Antonelli and Viera, 2015; Robinson,
38 Smith and Jones, 2021). Furthermore, a previous study indicates that colour-coding plays an
39 important role in people's decision-making processes. Colour-coding, in particular, may
40 improve analytical thinking rather than the quick and emotional thinking process, leading to
41 healthier choices (Cecchini and Warin, 2016; Fenko, Nicolaas, and Galetzka, 2018). While
42 many studies have been conducted on the use of colour-coding on front-of-pack labelling
43 (e.g., Sacks *et al.*, 2011; Mhurchu *et al.*, 2018; Fenko, Nicolaas, and Galetzka, 2018; Erdem and
44 Campbell, 2019; Hagmann and Siegrist, 2020), the use on menu labelling has gone unnoticed.
45
46
47
48
49
50

51 **3. Materials and Method**

52 **3.1. Study design**

53 We devised a field experiment in a natural eating environment at a restaurant in Stirling to
54 investigate the effect of restaurant menu labelling on consumers' meal choices and attitudes
55 toward the use of nutritional labels on menus. The restaurant is located in a hotel on campus
56 and is open to the public and students seven days a week. The experiment consisted of a
57 control condition in which no menu labels were included on the restaurant menu and a
58
59
60

1
2
3 treatment condition in which nutritional labels were included on the same menu used in the
4 control group. The labels were placed next to the menu items and contained information
5 about the dish's calorie and nutritional content (see Figure 1). The chefs provided us with the
6 recipes they used during the experiment, and we calculated the calorie content of each dish
7 using McCane and Widdowson's Composition of Foods Integrated Dataset (2006). This is a
8 database that is frequently used in calorie calculations and on which the MenuCal software is
9 based. We cross-validated our calculations with an expert dietitian at the company that
10 provided catering services to the restaurant after determining the calories and nutritional
11 contents of the restaurant's starters, main dishes, and desserts during the study period. We
12 designed new menu labels after calculating the calories and nutritional content of each dish.
13
14
15
16
17

18 In accordance with the standard practice used in front-of-pack labelling, the menu labels
19 included information about the calorie content of the entire portion and 100 grams of the
20 dish (see Figure 1). The labels also included information about the dishes' fat, saturated fat,
21 sugars, and salt content. We communicated nutrient content using traffic light coding and
22 qualitative descriptors for their levels, such as low, medium, and high, as found on food
23 product front-of-pack labelling. These classifications are based on the guidelines for
24 developing front-of-pack nutritional labelling for pre-packaged foods sold in retail outlets
25 (Department of Health, 2016). For example, if an item is coded with green, it indicates a low
26 level of that nutrient in 100 grams of that dish; amber indicates a medium level of nutrients
27 in 100 grams of the dish; and red indicates a high level of nutrients in 100 grams of the dish.
28 On the labels, the total portion size of the dish was also specified.
29
30
31
32
33

34 FIGURE 1 HERE

35 Another essential feature of the menu labels we created was how we communicated the
36 calories per 100 grams of food to the participants. We included the average amount of
37 walking time required to burn calories from the dish's consumption. The walking time is
38 calculated using the data from the BUPA calorie calculator¹ and cross-checked against other
39 sources such as the British Heart Foundation² and the NHS³. It is based on an adult who weighs
40 75 kg and walks at a moderate pace (3 miles per hour). Appendix A contains the menu with
41 labels (intervention). The menu in the control condition is identical to the one in the
42 treatment condition, but it lacks nutritional labels.
43
44
45
46

47 The experiment began by giving participants either the standard menu (control) or the same
48 menu with labels (treatment), depending on whether they were in the control or treatment
49 group. Following their meal, the waiting staff handed them a postcard with five quick
50 questions about their previous experience with nutritional labelling, perceived usefulness and
51 influence of nutritional labels on food choices made, gender, and food items ordered from
52 the menu. We were limited to using postcards with short questions due to the restaurateur's
53
54
55

56
57 ¹ <https://www.bupa.co.uk/health-information/tools-calculators/calories-calculator>

58 ² <https://www.bhf.org.uk/informationsupport/support/healthy-living/staying-active/exercise-calorie-calculator>

59 ³ <https://www.nhs.uk/live-well/healthy-weight/understanding-calories/>
60

1
2
3 concern about influencing their customers' dining experience with a standard follow-up
4 questionnaire. As a result, we were unable to investigate other factors that could have aided
5 in the investigation of food choice in greater depth.
6
7

8 We asked the control group if they were *familiar with nutritional labelling for foods*, if they
9 *would find nutritional information useful when selecting meals from a menu*, and if *this*
10 *information would influence what they chose*. The response options for these questions
11 ranged from "very familiar/very much agree" to "very unfamiliar/very much disagree" on a
12 five-point Likert scale. In addition to these three opinion questions, we asked participants
13 about their gender and the meals they ordered (starter, main, and dessert).
14
15

16 We asked the treatment group similar questions, but because the restaurant menu included
17 nutritional labels, our questions reflected this and sought more details on their opinions. We
18 inquired about their level of *familiarity with the labelling prior to visiting the restaurant*,
19 *whether the nutritional labelling influenced what they ordered, the most and least useful*
20 *information on the label*, their gender, and the meals they chose. While the response options
21 for label familiarity and influence were mirroring those in the control questionnaire, the
22 usefulness of labelling information included *calories, walking minutes, fat, saturates, sugars,*
23 *and salt contents*.
24
25
26

27 The items in the mini-questionnaires were chosen based on our review of the literature and
28 discussions with the restaurateur and local policymakers. Appendix B contains a copy of the
29 mini-questionnaires for the control and intervention groups. The University's Ethics Panel
30 approved the study protocol. Furthermore, before they were fielded, the mini questionnaires
31 were piloted with a small sample from the campus.
32
33
34

35 3.2. Data

36 In 2017, we collected data from 197 consumers. Participants in the study were neither
37 recruited nor chosen. They were restaurant patrons who chose to eat there on the spur of
38 the moment, unaware of the experiment. The treatment group had 108 participants,
39 compared to 89 in the control group. From March to July, we began collecting data from the
40 control group. The new menus were then labelled, and treatment data was collected from
41 July to December.
42
43
44

45 The participant profiles in the control and intervention groups are very similar. As shown in
46 Table 1, the proportion of males in each group is similar (45% control, 44% treatment), with
47 more people in the control group not reporting their gender and a slightly higher proportion
48 of females in the treatment group. Statistically speaking, there are no gender differences
49 between the control and treatment groups (Pearson $\chi^2(2) = 0.50$ with p-value = 0.77).
50
51
52
53

54 TABLE 1 HERE

55 It is critical for these types of natural experiments to have similar demographics for each
56 group in order to control for any unobservable variable biases within the analysis. There is a
57
58
59
60

1
2
3 slight difference in the male-to-female ratio between the groups, but this is not unusual given
4 the small sample size.
5

6 3.3. Analysis

7
8 The empirical evidence presented in the paper is based on descriptive and inferential
9 statistical analyses to compare and test differences between and within the control and
10 treatment conditions, and ordered logistic regression to investigate what predicts the
11 likelihood of labels perceived influential on food choices.
12
13

14 The inferential analysis is performed using Pearson's χ^2 to determine whether familiarity with
15 nutritional labelling, perception of the usefulness of menu labels, and the influence of labels
16 on food choices differ significantly between and within groups.
17
18

19 The ordered logistic regression allows us to predict the probability of labels as influential on
20 food choices using the perceived usefulness of calorie labels, familiarity with them, and
21 gender. Statistically, we can present the model as follows:
22

$$23 \log \frac{P(Y \leq j)}{P(Y \geq j)} = \beta_j + \beta X \quad j \in [1, J - 1], \quad \text{eq 1}$$

24
25
26
27 where $j \in [1, J - 1]$ are the levels of the ordinal outcome variable Y , agreement on the
28 influence of nutritional labels on food choices [disagree, neutral, agree]⁴. This outcome
29 variable is derived from a question on the control and treatment questionnaires that asks
30 participants' opinions on the influence of nutritional information on their food choices on a
31 five-point Likert-based scale⁵. The ordinal outcomes are distinguished by the $J - 1$
32 intercepts β_j . The benchmark level is J . β s are coefficients for X covariates: gender,
33 perceived usefulness of the labels and familiarity with the nutritional labelling.
34
35
36

37 By a simple algebraic manipulation, we can formulate the probability of observing an outcome
38 at a level j by:
39
40
41

$$42 p(Y \leq j) = \frac{\exp(\beta_j + \beta X)}{1 + \exp(\beta_j + \beta X)} \quad \text{eq 2}$$

43
44
45 This probability expression in eq 2 takes the following forms for the control (p_{ctrl}) and
46 treatment (p_{tr}):
47
48
49
50
51

52
53 ⁴ We note a slight difference in the meaning of the dependant variable for the control and treatment conditions.
54 The response to "influence" question in the control condition refers to how information on labels *would*
55 influence participant's food choice if they were provided. However, the treatment group made their food
56 choices using menus with labels, and therefore, we asked them whether information on labels *actually*
57 influenced their food choices.

58 ⁵ Due to the small sample size, few responses were given to each level of Likert scale. For this reason, we decided
59 to derive a variable with three levels, rather than using all five levels.
60

$$p_{ctrl} = \frac{\exp(\beta_j + \beta_1 X_{female} + \beta_2 - 3X_{familiar} + \beta_4 - 5X_{useful})}{1 + \exp(\beta_j + \beta_1 X_{female} + \beta_2 - 3X_{familiar} + \beta_4 - 5X_{useful})} \quad \text{eq 3}$$

$$p_{tr} = \frac{\exp(\beta_j + \beta_1 X_{female} + \beta_2 - 3X_{familiar} + \beta_4 X_{usefulcal} + \beta_5 X_{usefulwalk})}{1 + \exp(\beta_j + \beta_1 X_{female} + \beta_2 - 3X_{familiar} + \beta_4 X_{usefulcal} + \beta_5 X_{usefulwalk})} \quad \text{eq 4}$$

where female is defined as a binary variable, familiarity is a categorical variable with three levels (unfamiliar, neutral, familiar), "neutral" being the baseline level. The "usefulness" variables are defined differently in control and treatment. It asks the level of agreement on whether nutritional labels are useful on food choices in the control condition (disagree, neutral, agree); in the treatment condition, it asks the most and least useful information on nutritional labels. We created two dummy variables from the treatment questionnaire that were related to the usefulness of calorie ("usefulcal") and walking minute ("usefulwalk").

We then use maximum likelihood estimation to estimate the β coefficients that maximise the log-likelihood function.

$$\sum_{i=1}^N [Y_i * \ln(P_{Y_i}) + (1 - Y_i) * \ln(1 - P_{Y_i})], \quad \text{eq 5}$$

where N is the number of participants.

4. Findings

4.2. Exploratory Data Analysis:

The data provided information on the gender of the participants, their familiarity with nutritional labels, their opinions on how much the menu labels influence their food choices, and their actual menu choices. This section presents descriptive statistics and the results of statistical tests performed between and within groups, as shown in Table 1.

TABLE 1 HERE

Our exploratory analysis shows that control and treatment samples are not statistically significantly different in gender distribution (Pearson $\chi^2(2) = 3.71$, p-value = 0.15). There are, however, some significant differences in terms of familiarity with the nutritional labels (Pearson $\chi^2(4) = 7.67$, p-value = 0.10) and the perceived influence of labels on actual food choices from the menu (Pearson $\chi^2(4) = 12.81$, p-value = 0.01). We see that the control has a slightly higher proportion of individuals more familiar with nutritional labelling (68%) than individuals in the treatment group (56%). When we further investigate the familiarity within each sample, we see no significant association with gender (control: Pearson $\chi^2(8) = 10.99$, p-value = 0.20; treatment: Pearson $\chi^2(8) = 7.52$, p-value = 0.48). However, we see a significant association between the perceived usefulness and familiarity with labels in the control condition (Pearson $\chi^2(8) = 17.72$, p-value = 0.001).

1
2
3 Regarding the influence the nutritional labels have on individuals' meal choices, we find
4 significant differences in views between the control and the treatment groups (Pearson $\chi^2(4)$
5 = 12.81 with p-value = 0.01). While 23% of the control group disagreed that the nutritional
6 labelling influenced what they chose from the menu, this was 43% for the treatment group,
7 who were actually presented with menus including labels. Nearly equal proportions of the
8 samples agreed (c.40%) that nutritional labels influenced their choices. This indicates that
9 more people are neutral in their opinion in the control group than in the treatment. This could
10 be due to the uncertainty or unfamiliarity about how actual nutritional labels would look like
11 on restaurant menus. When we reviewed the feedback participants left over the survey card,
12 a few participants mentioned that the nutritional labels on the menu "spoiled their dining
13 experience" as they did not want to know the nutritional details of their meals (a
14 phenomenon called strategic self-ignorance (Thunström *et al.*, 2016)). These findings are
15 consistent with other studies that found a similar proportion of individuals would not use
16 nutritional labels on menus or not find them influential (Burton *et al.*, 2006; Harnack and
17 French, 2008; Liu *et al.*, 2012; McCann *et al.*, 2013; Nikolaou, Lean and Hankey, 2014).

18
19 Looking at which information on the label is viewed as the most influential in the treatment
20 group's decision-making process, we find that both calorie information and walking minutes
21 to burn calories are most influential in making choices, as shown in Figure 2 below. Nutrient
22 information (such as salt, fat, saturates, and sugar) ranked low in terms of usefulness in
23 decision-making. Participants who believe nutrients are the most useful information on labels
24 also believe walking minutes are the least useful. This could be due to a lack of familiarity with
25 the measure. When we explore if the usefulness of certain information on labels varies by
26 gender, we find no significant differences.

27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURE 2 HERE

Using the data provided by the survey respondents⁶, we calculate the average calories ordered by the control and treatment group for each course. As shown in Table 1, the differences between average calories for starter, main dish, and desserts are very small and statistically insignificant, as evident from the result of the t-tests. Furthermore, although the average total calories in the treatment group are less than the average total calories in the control group, this difference is not statistically significant, either ($t(170) = 0.36, p = 0.72$). This is consistent with findings from Cawley, Susskind and Willage (2020) study on calorie labelling of restaurant menus.

4.2. Regression Analysis

We conducted an ordered logistic regression to determine how nutritional labelling of menus affects actual food choices and what predicts the probability of labels being influential on meal choice. The dependent variable is an ordinal variable with three levels: disagreement, neutral opinion, and agreement on the influence of nutritional labels on food choices⁷.

⁶ Not all participants provided details on what they ordered from the menu.

⁷ We explored various regression models, such as multinomial logit model using food choices as outcome variable, OLS using total calorie consumption as the outcome variable, and logistic regression using a binary

1
2
3 The usefulness of nutritional labels was measured on a Likert-based scale, and we recoded it
4 in this regression analysis using three ordinal levels: disagreeing on the usefulness of labels,
5 neutral, and agreeing on the usefulness of labels. However, because the treatment groups
6 had already encountered the nutritional labels during the experiment, we asked them which
7 aspects (i.e., calorie content, walking minutes, nutrient contents) they found most useful and
8 which they found least useful. Because of this nuance, there is no direct comparison of the
9 perceived usefulness of the labels between control and treatment.
10
11

12
13 According to the results, familiarity with nutritional labels has no significant effect on the
14 labels' likelihood of influencing food choices in both control and treatment groups. On the
15 other hand, the perceived usefulness of the labels has some significant effect on the likelihood
16 of labels being influential during food choices.
17
18

19 In the control group, keeping all variables constant, if participants agree with the usefulness
20 of the labels, they are 3.29 (odds ratio) times more likely to find the nutritional labels
21 influential on their meal choices if they were provided with a menu with nutritional labels
22 (relative to the participants who neither agree nor disagree with it). In the treatment group,
23 if calorie information is believed to be the most useful aspect of the menu label, participants
24 are 3.51 (odd ratio) times more likely to find the nutritional labels influential when making
25 meal choices from a menu with labels (compared to a case where nutrient contents are
26 considered as the most useful aspect). Walking minutes, on the other hand, is not statistically
27 significant in predicting the likelihood of influence on food choices (compared to nutrient
28 details).
29
30
31

32
33 In terms of gender, females in the control group are 3.34 times (odds ratio) more likely to find
34 nutritional labels persuasive. However, in the treatment group, we found no significant
35 gender effect on the likelihood of nutritional labels being viewed as important on food
36 choices.
37
38

39 TABLE 2 HERE
40

41 5. Discussion

42
43 This research used a field experiment at a restaurant to investigate the effect of nutritional
44 labelling on restaurant menus on actual meal purchases in a real-world setting. While it does
45 not seek to add conclusive weight to the policy debate, it does provide empirical evidence on
46 the impact of menu labels on meal choices at restaurants and how consumers perceive them.
47 Several studies on nutritional labelling have had equivocal results (Restrepo, 2017; S. Bleich
48 *et al.*, 2017; S. N. Bleich *et al.*, 2017; Kerins *et al.*, 2018). This paper also did not find conclusive
49 evidence that menu calorie labelling is effective. This discovery can be attributed to a number
50 of factors. To begin with, while field experiments are useful for gathering real-life data
51 without influencing participants and so have higher ecological validity than lab-based studies,
52 they also have certain flaws that may explain our rather poor results. These include (i) taking
53
54
55
56
57

58 outcome variable representing if calorie consumption is over the sample mean or median calorie consumption.
59 However, we did not find significant effects to report (these are available from the author upon request).
60

1
2
3 more time to collect data, which reduces the sample size of a study; (ii) having no control over
4 variables, which causes replicability and generalisability issues (external validity); and (iii)
5 having no control over attrition, which results in unbalanced/missing data in the analysis. The
6 amount of data collected is also determined by the restaurateur's desire to provide room
7 for data gathering after their customers' eating experience. In our situation, we agreed with
8 the company to ask a modest number of opinion questions to elicit the important features of
9 menu labelling use. As a result, the lack of rich data was not surprising, as many systematic
10 reviews in the literature have noted (Sinclair, Cooper, and Mansfield, 2014; Arno and Thomas,
11 2016; Crockett *et al.*, 2018). Despite these limitations, this paper is useful for researchers and
12 others who want to design a study investigating the impact of menu labelling on food choices
13 in a real-world setting.
14
15
16
17

18 One of the goals of this study is to demonstrate the importance of information disclosure on
19 restaurant menus. We discovered that disclosing nutritional and calorie content information
20 on restaurant menus increased participants' perceptions of labels as influential on food
21 choices. There is evidence that menu labelling information can have varying levels of
22 importance to consumers, and the more useful the information is to consumers, the more
23 likely they will be influenced by it (Breck *et al.*, 2014). Because calories are useful to some
24 people, healthy choices are often made when people are knowledgeable about calories and
25 can use the information on labels (Girz *et al.*, 2012; Ni Mhurchu *et al.*, 2018). We wanted to
26 show that visual cues like pictograms and colour coding would result in more easily accessible
27 and, as a result, effective menu labelling in our study. We discovered that a higher proportion
28 of people found the pictogram to be the most useful part of the menu labelling. This is
29 because of the visual presentation with a familiar activity aids in translating and thus
30 understanding numeric calorie information. Including contextual information, such as colour
31 coding or infographics, on menu labels, according to studies, is more effective than just
32 displaying calorie content (Liu *et al.*, 2012; Morley *et al.*, 2013; Antonelli and Viera, 2015).
33
34
35
36
37
38

39 However, when the actual calories of meals ordered are compared to perceived influence,
40 there are no significant differences between the control and treatment groups. This indicated
41 that participants in the treatment condition believed nutritional labels had an impact on their
42 decisions; however, their actual purchased calories were not statistically significantly lower
43 than those in the control conditions. We should point out that we only observed the orders
44 and not the consumption. As a result, it is possible that participants respond to nutritional
45 labels by changing the amount they consume rather than changing their orders.
46 Unfortunately, we cannot prove this claim, but it would be useful if future studies could test
47 how labels influence consumption.
48
49
50

51 Females are more likely than males to be familiar with nutritional menu labels, which is
52 consistent with previous studies (Breck *et al.*, 2014; Nikolaou, Lean, and Hankey, 2014).
53 However, a higher level of familiarity does not always imply comprehension or use of these
54 labels.
55
56

57 Our findings indicate that the "perceived" usefulness of labels appears to be related to
58 whether labels are thought to influence meal choices. This is a significant finding. As Spinks
59 and Mortimer (2015) point out, the use of pictograms and colour-coded nutritional
60

1
2
3 information, including calorie content, may be some ways to contextualise difficult-to-
4 understand information on menu labels, and thus, might be influencing the perceived
5 “usefulness” of labels.
6
7

8 While most studies on menu labelling have focused on the consumer's point of view (Din,
9 Zahari, and Shariff, 2012; Restrepo, 2017), a few studies have looked at business impacts
10 (Fitzgerald *et al.*, 2018). While this study focuses on consumer responses to menu labelling,
11 we collected informal feedback from the restaurateur during the field experiment's design
12 and implementation. Many of the barriers to menu labelling implementation identified by
13 restaurateurs have been extensively discussed in the literature. For example, measuring
14 calories was difficult due to a lack of knowledge, expertise, and equipment. This problem was
15 exacerbated by variations in cooking methods and recipe changes that altered the calorie and
16 nutritional content of meals. While most of the barriers were related to implementing menu
17 labelling and how it affected how the kitchen ran, potentially stifling chef experimentation
18 and increasing difficulty infrequently changing menu items (Obbagy *et al.*, 2011; Condrasky
19 *et al.*, 2015), which could lead to potential profit loss. There was also concern that, in order
20 to implement menu labelling, restaurant staff would need to be trained and able to answer
21 questions about these labels, similar to findings in other studies (e.g., Thomas, 2015). This
22 could be difficult due to a lack of resources and time for training, as well as potential issues
23 with waiting staff turnover.
24
25
26
27
28

29 While there were concerns and barriers to implementing menu labelling at the restaurant,
30 there was a genuine desire to provide customers with more information to help them make
31 healthy choices. However, this information had to be balanced against the risk of
32 overwhelming the customer and negatively impacting their dining experience (Hodgkins *et*
33 *al.*, 2012).
34
35

36 Another interesting observation from this study was that the restaurant decided to
37 reformulate a few items on their menu when designing the labels. This involved reducing the
38 portion sizes to have a lower level of fat and sugar content. This reformulation is often
39 credited as a potential benefit of menu labelling (Bleich *et al.*, 2017; Vandevijvere and
40 Vanderlee, 2019).
41
42
43

44 There are several important directions for future research. First, field experiments should be
45 conducted in restaurants varying in size, location, customer base, and menu offerings.
46 Second, the actual consumption data should also be collected to test the effect of menu labels
47 on meal consumption.
48
49

50 **6. Concluding remarks**

51 This paper presents empirical evidence on the effect of menu labelling on actual meal
52 purchases by consumers in a real-world setting. While the findings do not add conclusive
53 weight to the policy debate, they do add to the literature by providing empirical evidence
54 from a field experiment and sharing experiences from the study's design and implementation.
55 More research is needed, however, to address the limitations highlighted in this study. The
56 requirement for useful menu labelling necessitates the testing of various labelling
57
58
59
60

1
2
3 interventions and formats. A more in-depth study could reveal whether different menu
4 labelling formats can reduce calorie purchase, as well as consumption. However, more
5 engagement with businesses of various sizes and sectors of the restaurant industry is required
6 before any policy recommendation on menu labelling implementation can be made.
7
8

9 **Acknowledgements**

10 To be added.
11
12

13 **References**

14
15 Alexander, M., O'Gorman, K. D. and Woods, K. (2010) 'Nutritional labelling in restaurants:
16 whose responsibility is it anyway? International', *International Journal of Contemporary*
17 *Hospitality*, 22(4), pp. 572–579.
18

19
20 Almanza, B. A., Nelson, D. and Chai, S. (1997) 'Obstacles to nutrition labelling in restaurants',
21 *Journal of the American Dietetic Association*, pp. 157–161. doi: 10.1016/S0002-
22 8223(97)00041-2.
23

24
25 Antonelli, R. and Viera, A. J. (2015) 'Potential effect of physical activity calorie equivalent
26 (PACE) labeling on adult fast food ordering and exercise', *PLoS ONE*, 10(7). doi:
27 10.1371/journal.pone.0134289.

28
29 Arno, A. and Thomas, S. (2016) 'The efficacy of nudge theory strategies in influencing adult
30 dietary behaviour: A systematic review and meta-analysis', *BMC Public Health*. BioMed
31 Central Ltd., 16(1). doi: 10.1186/s12889-016-3272-x.

32
33 Auchincloss, A. H. *et al.* (2013) 'Barriers and facilitators of consumer use of nutrition labels at
34 sit-down restaurant chains', *Public Health Nutrition*, 16(12), pp. 2138–2145. doi:
35 10.1017/S1368980013000104.
36

37
38 Auchincloss, A. H. *et al.* (2014) 'Nutritional value of meals at full-service restaurant chains',
39 *Journal of Nutrition Education and Behavior*. Elsevier Inc., 46(1), pp. 75–81. doi:
40 10.1016/j.jneb.2013.10.008.

41
42 Bassett, M. T. *et al.* (2008) 'Purchasing behavior and calorie information at fast-food chains in
43 New York City, 2007', *American Journal of Public Health*, 98(8), pp. 1457–1459. doi:
44 10.2105/AJPH.2008.135020.

45
46 Bleich, S. *et al.* (2017) 'Calorie changes in large chain restaurants from 2008 to 2015',
47 *Preventative medicine*, 199, p. 1120116. Available at:
48 <https://www.sciencedirect.com/science/article/pii/S009174351730124X> (Accessed: 21
49 October 2019).
50

51
52 Bleich, S. N. *et al.* (2017) 'A Systematic Review of Calorie Labeling and Modified Calorie
53 Labeling Interventions: Impact on Consumer and Restaurant Behavior', *Obesity*. Blackwell
54 Publishing Inc., pp. 2018–2044. doi: 10.1002/oby.21940.

55
56 Bleich, S. N., Moran, A. J. and Jarlenski, M. P. (2018) 'Higher-calorie menu items eliminated in
57 large chain restaurants', *American Journal of Preventive Medicine*, 54(2), pp. 214–220.
58 Available at: <https://www.sciencedirect.com/science/article/pii/S0749379717306669>
59 (Accessed: 4 November 2019).
60

1
2
3
4
5
6
7
8
9
Bleich, S. N., Wolfson, J. A. and Jarlenski, M. P. (2016) 'Calorie changes in large chain restaurants: declines in new menu items but room for improvement', *American Journal of Preventive Medicine*, 50(1), pp. 1–8. Available at: <https://www.sciencedirect.com/science/article/pii/S0749379715002305> (Accessed: 4 November 2019).

10
11
12
Bollinger, B., Leslie, P. and Sorensen, A. (2011) 'Calorie posting in chain restaurants', *American Economic Journal: Economic Policy*, 3(1), pp. 91–128. doi: 10.1257/pol.3.1.91.

13
14
15
Breck, A. *et al.* (2014) 'Who reports noticing and using calorie information posted on fast food restaurant menus?', *Appetite*. Elsevier Ltd, 81, pp. 30–36. doi: 10.1016/j.appet.2014.05.027.

16
17
18
Britt, J. W. *et al.* (2011) 'Feasibility of Voluntary Menu Labeling Among Locally Owned Restaurants', *Health Promotion Practice*, 12(1), pp. 18–24. doi: 10.1177/1524839910386182.

19
20
21
22
23
24
25
Bruemmer, B. *et al.* (2012) 'Energy, saturated fat, and sodium were lower in entrées at chain restaurants at 18 months compared with 6 months following the implementation of mandatory', *Journal of the Academy of Nutrition and Dietetics*, 112(8), pp. 1169–1176. Available at: <https://www.sciencedirect.com/science/article/pii/S2212267212005928> (Accessed: 21 October 2019).

26
27
28
29
Burton, S., Creyer, E. H., Kees, J., & Huggins, K. (2006). Attacking the obesity epidemic: the potential health benefits of providing nutrition information in restaurants. *American journal of public health*, 96(9), 1669-1675.

30
31
32
33
34
Cawley, J., Susskind, A., & Willage, B. (2020). The Impact of information disclosure on consumer behavior: Evidence from a randomised field experiment of calorie labels on restaurant menus. *Journal of Policy Analysis and Management*, 39(4), 1020-1042.

35
36
37
38
39
Cecchini, M., & Warin, L. (2016). Impact of food labelling systems on food choices and eating behaviours: a systematic review and meta-analysis of randomised studies. *Obesity reviews: an official journal of the International Association for the Study of Obesity*, 17(3), 201–210. <https://doi.org/10.1111/obr.12364>.

40
41
42
43
44
Condrasky, M. D. *et al.* (2015) 'Opinions, Knowledge, and Current Practices of Culinary Arts Instructors and Professionals in Regards to Healthy Food Techniques', *Journal of Culinary Science and Technology*, 13(4), pp. 287–302. doi: 10.1080/15428052.2015.1015669.

45
46
47
48
Crockett, R. A. *et al.* (2018) 'Nutritional labelling for healthier food or non-alcoholic drink purchasing and consumption', *Cochrane Database of Systematic Reviews*. John Wiley and Sons Ltd. doi: 10.1002/14651858.CD009315.pub2.

49
50
51
Department of Health. <https://www.food.gov.uk/business-guidance/nutrition-labelling> (Accessed: 1st of January 2017).

52
53
54
55
Din, N., Zahari, M. S. M. and Shariff, S. M. (2012) 'Customer Perception on Nutritional Information in Restaurant Menu', *Procedia - Social and Behavioral Sciences*, 42(July 2010), pp. 413–421. doi: 10.1016/j.sbspro.2012.04.205.

56
57
58
59
60
Ellison, B., Lusk, J. L. and Davis, D. (2013) 'Looking at the label and beyond: the effects of calorie labels, health consciousness, and demographics on caloric intake in restaurants', *International Journal of Behavioral Nutrition and Physical Activity*, 10(1), p. 21. doi: 10.1186/1479-5868-10-21.

1
2
3 Erdem, S. and Campbell, D. (2019) 'Investigating System 1 and 2 Decision-making in a Food
4 Choice and Eye Tracking Experiment', in International Choice Modelling Conference. Tokyo,
5 Japan.
6

7 Fenko, A., Nicolaas, I., & Galetzka, M. (2018). Does attention to health labels predict a healthy
8 food choice? An eye-tracking study. *Food quality and preference*, 69, 57-65.
9

10 Food Safety Authority of Ireland (2012) 'Putting calories on menus in Ireland-Draft technical
11 guidance for food businesses'.
12

13 Food Standards Scotland (2018) *FSS Response to Cochrane Review of Nutrition Labelling on*
14 *Menus | Food Standards Scotland, Food Standards Scotland Website*. Available at:
15 [https://www.foodstandards.gov.scot/news-and-alerts/fss-response-to-cochrane-review-of-](https://www.foodstandards.gov.scot/news-and-alerts/fss-response-to-cochrane-review-of-nutrition-labelling-on-menus)
16 [nutrition-labelling-on-menus](https://www.foodstandards.gov.scot/news-and-alerts/fss-response-to-cochrane-review-of-nutrition-labelling-on-menus) (Accessed: 21 October 2019).
17
18

19 Geaney, F. *et al.* (2015) *Evaluation of the uptake of voluntary calorie posting on menus in*
20 *Ireland*.
21

22 Girz, L. *et al.* (2012) 'The effects of calorie information on food selection and intake',
23 *International Journal of Obesity*, 36(10), pp. 1340–1345. doi: 10.1038/ijo.2011.135.
24

25 Hagmann, D., & Siegrist, M. (2020). Nutri-Score, multiple traffic light and incomplete
26 nutrition labelling on food packages: Effects on consumers' accuracy in identifying healthier
27 snack options. *Food Quality and Preference*, 83, 103894.
28
29

30 Hammond, D. *et al.* (2013) 'A randomised trial of calorie labeling on menus', *Preventive*
31 *Medicine*. Elsevier Inc., 57(6), pp. 860–866. doi: 10.1016/j.ypmed.2013.09.020.
32

33 Harnack, L. J. and French, S. A. (2008) 'Effect of point-of-purchase calorie labeling on
34 restaurant and cafeteria food choices: A review of the literature', *International Journal of*
35 *Behavioral Nutrition and Physical Activity*, 5, pp. 3–8. doi: 10.1186/1479-5868-5-51.
36

37 Hodgkins, C. *et al.* (2012) 'Understanding how consumers categorise nutritional labels: A
38 consumer derived typology for front-of-pack nutrition labelling', *Appetite*. Elsevier Ltd, 59(3),
39 pp. 806–817. doi: 10.1016/j.appet.2012.08.014.
40
41

42 Holmes, A. S. *et al.* (2013) 'Effect of different children's menu labeling designs on family
43 purchases', *Appetite*, 62, pp. 198–202. doi: 10.1016/j.appet.2012.05.029.
44

45 Kelly, S. M. *et al.* (2014) 'Putting calories on the menu in Ireland: evaluation of an online
46 calorie calculator for food businesses', *Proceedings of the Nutrition Society*, 73(OCE2), p.
47 11419. doi: 10.1017/s0029665114000883.
48

49 Kerins, C. *et al.* (2018) 'Barriers and facilitators to implementation of menu labelling
50 interventions to support healthy food choices: A mixed methods systematic review protocol',
51 *Systematic Reviews*. BioMed Central Ltd., 7(1). doi: 10.1186/s13643-018-0752-3.
52

53 Knai, C. *et al.* (2015) 'Has a public-private partnership resulted in action on healthier diets in
54 England? An analysis of the Public Health Responsibility Deal food pledges', *Food Policy*.
55 Elsevier Ltd, 54, pp. 1–10. doi: 10.1016/j.foodpol.2015.04.002.
56
57

58 Kuo, T. *et al.* (2009) 'Menu labeling as a potential strategy for combating the obesity epidemic:
59 A health impact assessment', *American Journal of Public Health*, 99(9), pp. 1680–1686. doi:
60

1
2
3 10.2105/AJPH.2008.153023.
4

5 Liu, P. J. *et al.* (2012) 'A test of different menu labeling presentations', *Appetite*. Elsevier Ltd,
6 59(3), pp. 770–777. doi: 10.1016/j.appet.2012.08.011.
7

8 Mackison, D., Wrieden, W. and Anderson, A. (2009) 'Making an informed choice in the
9 catering environment: What do consumers want to know?', *Journal of Human Nutrition and*
10 *Dietetics*, 22(6), pp. 567–573. doi: 10.1111/j.1365-277X.2009.01000.x.
11

12 Mah, C. L. *et al.* (2013) 'Ready for policy? stakeholder attitudes toward menu labelling in
13 Toronto, Canada', *Canadian Journal of Public Health*, 104(3). doi: 10.17269/cjph.104.3708.
14

15 McCann, M. T. *et al.* (2013) 'Influence of nutrition labelling on food portion size consumption',
16 *Appetite*. Elsevier Ltd, 65, pp. 153–158. doi: 10.1016/j.appet.2013.02.013.
17

18 Mhurchu, C. N., Eyles, H., Jiang, Y., & Blakely, T. (2018). Do nutrition labels influence
19 healthier food choices? Analysis of label viewing behaviour and subsequent food purchases
20 in a labelling intervention trial. *Appetite*, 121, 360-365.
21
22

23 Morley, B. *et al.* (2013) 'What types of nutrition menu labelling lead consumers to select less
24 energy-dense fast food? An experimental study', *Appetite*. Elsevier Ltd, 67, pp. 8–15. doi:
25 10.1016/j.appet.2013.03.003.
26

27 Obbagy, J. E. *et al.* (2011) 'Chefs' opinions about reducing the calorie content of menu items
28 in restaurants', *obesity*. Nature Publishing Group, 19(2), pp. 332–337. doi:
29 10.1038/oby.2010.188.
30
31

32 Orfanos, P. *et al.* (2009) 'Eating out of home: Energy, macro-and micronutrient intakes in 10
33 european countries. The European prospective investigation into cancer and nutrition',
34 *European Journal of Clinical Nutrition*, 63, pp. S239–S262. doi: 10.1038/ejcn.2009.84.
35

36 Pomeranz, J. L. (2011) 'Front-of-package food and beverage labeling: New directions for
37 research and regulation', *American Journal of Preventive Medicine*. Elsevier Inc., 40(3), pp.
38 382–385. doi: 10.1016/j.amepre.2010.11.009.
39

40 Restrepo, B. J. (2017) 'Calorie Labeling in Chain Restaurants and Body Weight: Evidence from
41 New York', *Health Economics (United Kingdom)*. John Wiley and Sons Ltd, 26(10), pp. 1191–
42 1209. doi: 10.1002/hec.3389.
43
44

45 Reale, S., Flint, S. and Capehorn, M. (2015) 'The impact of menu labelling on food choice in an
46 obese population', *Appetite*. Elsevier BV, 87, p. 391. doi: 10.1016/j.appet.2014.12.164.
47

48 Sacks, G., Tikellis, K., Millar, L., & Swinburn, B. (2011). Impact of 'traffic-light' nutrition
49 information on online food purchases in Australia. *Australian and New Zealand journal of*
50 *public health*, 35(2), 122-126.
51

52 Spinks, J. and Mortimer, D. (2015) 'The effect of traffic lights and regulatory statements on
53 the choice between complementary and conventional medicines in Australia: Results from a
54 discrete choice experiment', *Social Science and Medicine*. Elsevier Ltd, 124, pp. 257–265. doi:
55 10.1016/j.socscimed.2014.11.049.
56
57

58 Thomas, E. (2015) 'Commentary Food for thought: obstacles to menu labelling in restaurants
59 and cafeterias'. doi: 10.1017/S1368980015002256.
60

1
2
3 Thunström, L., Nordström, J., Shogren, J. F., Ehmke, M., & van't Veld, K. (2016). Strategic self-
4 ignorance. *Journal of Risk and Uncertainty*, 52(2), 117-136.
5

6 Vandevijvere, S. and Vanderlee, L. (2019) 'Effect of Formulation, Labelling, and Taxation
7 Policies on the Nutritional Quality of the Food Supply', *Current Nutrition Reports*. Springer
8 Science and Business Media LLC, 8(3), pp. 240–249. doi: 10.1007/s13668-019-00289-x.
9

10 Wootan, M. G. and Osborn, M. (2006) 'Availability of nutrition information from chain
11 restaurants in the United States', *American Journal of Preventive Medicine*, 30(3), pp. 266–
12 268. doi: 10.1016/j.amepre.2005.10.006.
13

14 Wootan, M. G., Osborn, M. and Malloy, C. J. (2006) 'Availability of point-of-purchase nutrition
15 information at a fast-food restaurant', *Preventive Medicine*, 43(6), pp. 458–459. doi:
16 10.1016/j.ypmed.2006.07.008.
17

18 Yepes, M. F. (2015) 'Mobile Tablet Menus: Attractiveness and Impact of Nutrition Labeling
19 Formats on Millennials' Food Choices', *Cornell Hospitality Quarterly*, 56(1), pp. 58–67. doi:
20 10.1177/1938965514546371.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Results of exploratory data analysis

	Control	Treatment	Test Statistics
Gender			
<i>Male</i>	48 (54%)	48 (47%)	Pearson $\chi^2(2) = 3.71$ p-value = 0.15
<i>Female</i>	34 (39%)	47 (46%)	
<i>Prefer not to say</i>	6 (7%)	8 (7%)	
N	88	103	
Familiarity with labels			
<i>Very unfamiliar</i>	3 (3%)	13 (13%)	Pearson $\chi^2(4) = 7.67$ p-value = 0.10
<i>Somewhat unfamiliar</i>	14 (16%)	13 (13%)	
<i>Neutral</i>	11 (13%)	19 (18%)	
<i>Somewhat familiar</i>	39 (44%)	35 (34%)	
<i>Very familiar</i>	21 (24%)	22 (22%)	
<i>N/A</i>	0 (%)	1 (1%)	
N	88	103	
Influence of labels on food choice			
<i>Very much disagree</i>	9 (10%)	28 (27%)	Pearson $\chi^2(4) = 12.81$ p-value = 0.01
<i>Somewhat disagree</i>	11 (13%)	16 (16%)	
<i>Neutral</i>	29 (33%)	18 (17%)	
<i>Somewhat agree</i>	30 (34%)	29 (28%)	
<i>Very much agree</i>	8 (9%)	12 (12%)	
<i>N/A</i>	1 (1%)	0 (0%)	
N	88	103	
Average calories of foods purchased (Kcal)			
<i>Starter</i>	281 (sd = 122)	275 (sd = 141)	t(149) = 0.25, p = 0.80
<i>Main</i>	666 (sd = 145)	670 (sd = 125)	t(168) = -0.19, p = 0.85
<i>Dessert</i>	456 (sd = 175)	424 (sd = 167)	t(127) = 1.04, p = 0.30
<i>Total calories</i>	1242 (sd = 314)	1224 (sd = 319)	t(170) = 0.36, p = 0.72
N	70	102	

Table 2. Ordered logistic regression result

<i>Dep var = influence (ordinal)</i> (1=Disagree, 2 = Neutral, 3= Agree)	Control		Treatment	
	Coef.	Odd Ratio (OR)	Coef.	Odd Ratio (OR)
Coefficients:				
<i>Familiarity: unfamiliar</i>	0.52 (0.79)	1.68 (0.79)	0.30 (0.62)	1.35 (0.62)
<i>Familiarity: neutral</i>	Baseline			
<i>Familiarity: familiar</i>	-0.02 (0.72)	0.97 (0.72)	0.30 (0.54)	1.35 (0.54)
<i>Usefulness: disagree</i>	-2.63*** (0.77)	0.08*** (0.77)		
<i>Usefulness: neutral</i>	Baseline			
<i>Usefulness: agree</i>	1.19** (0.59)	3.29** (0.59)		
<i>Most useful: calories</i>			1.25** (0.56)	3.51** (0.56)
<i>Most useful: walking minutes</i>			0.77 (0.51)	2.17 (0.51)
<i>Most useful: nutrient contents</i>	Baseline			
Female	1.21*** (0.49)	3.34*** (0.49)	0.12 (0.42)	1.12 (0.42)
Intercepts:				
Disagree unsure	-1.07 (0.67)		0.59 (0.62)	
Unsure agree	1.23* (0.66)		1.31** (0.63)	
N	81		89	
Log-likelihood	-65.70		-88.45	
***, **, * indicate significance at the 1%, 5%, and 10% levels, respectively. Standard errors are in parentheses.				




Figure 1. An example of a nutritional label used in the experiment





Figure 2. Most useful information on the nutritional labelling

Appendix A: Labelled Menu


STARTERS

Seasonal soup of the day 4.25


Crispy Stornoway black pudding and confit pork sandwich
with celeriac slaw, truffled spinach,
caramelised apples and sage jus 6.75

Per 100g	
242 kCal	Fat MED
	Saturates HIGH
	Sugars LOW
	Salt MED
48 mins walk	
A typical dish is 220g (533Kcal)	


Hot oaked-roast Scottish salmon and herb mousse
avocado, lemon and shallot dressing 6.50

Per 100g	
148 kCal	Fat MED
	Saturates MED
	Sugars LOW
	Salt MED
30 mins walk	
A typical dish is 135g (200Kcal)	


Terrine of chicken liver and port parfait
house chutney, Earl Grey soaked sultanas, brioche bun 6.50

Per 100g	
258 kCal	Fat MED
	Saturates HIGH
	Sugars MED
	Salt MED
52 mins walk	
A typical dish is 127g (327Kcal)	

Crispy Inverloch Goat's Cheese
Balsamic, locally sourced honey-roasted garden beetroot,
rocket leaves 6.00

Per 100g	
281 kCal	Fat HIGH
	Saturates HIGH
	Sugars MED
	Salt MED
56 mins walk	
Typical dish is 105g (295Kcal)	

Crown of seasonal melon
exotic fruits and Arran mango sorbet 5.65


Per 100g	
37 kCal	Fat LOW
	Saturates LOW
	Sugars MED
	Salt LOW
7 mins walk	
A typical dish is 258g (95Kcal)	

Journal

MAIN COURSE


Roast rump of Perthshire lamb

braised lentils, button onions, pancetta, apple and elderflower chutney, wilted greens, madeira jus 16.95

Per 100g	
185 kCal	Fat MED
	Saturates MED
	Sugars LOW
	Salt MED
37 mins walk	
A typical dish is 345g (638Kcal)	


Poached smoked Scottish haddock

asparagus, leek and Arran mustard potatoes, soft poached egg, sauce hollandaise 14.95

Per 100g	
105 kCal	Fat MED
	Saturates MED
	Sugars LOW
	Salt MED
21 mins walk	
A typical dish is 375g (394Kcal)	


Slow cooked daube of highland beef

horseradish mash, wild mushrooms, fine beans, Bordelaise sauce 15.25

Per 100g	
157 kCal	Fat MED
	Saturates MED
	Sugars LOW
	Salt LOW
31 mins walk	
A typical dish is 350g (550Kcal)	


Grilled fillet of Scottish salmon

lemon and soft herb risotto, crisp Parma ham, pesto dressing 14.50

Per 100g	
267 kCal	Fat MED
	Saturates MED
	Sugars LOW
	Salt MED
53 mins walk	
A typical dish is 300g (801Kcal)	


Wild mushroom and asparagus filo parcel

spinach, braised rice and a sweet chilli dressing 13.50

Per 100g	
234 kCal	Fat MED
	Saturates MED
	Sugars LOW
	Salt MED
47 mins walk	
Typical dish is 303g (708Kcal)	

8oz Buccleuch chargrilled sirloin steak

fries, flat cap mushroom, roasted plum tomato, garlic and herb butter 19.95

Per 100g	
199 kCal	Fat MED
	Saturates HIGH
	Sugars LOW
	Salt MED
40 mins walk	
A typical dish is 422g (840Kcal)	

Chef's seasonal dish of the day 14.50

Ask your server for details

**Nutritional details and calories are average values for a typical dish. The walking time is for an average adult walking at a moderate pace.


Final

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

DESSERTS


Baked Bramley apple crumble

Arran vanilla bean ice cream, spiced red wine syrup 4.50




Lemon and meringue

lemon curd mousse, fluffy meringue, white chocolate sauce 5.95




Hazelnut delice

Arran peanut butter ice cream, dulce de leche 6.50




Classic chocolate fondant,

poached sour cherries, vanilla mascarpone, pistachio wafer 5.75



Selection of Scottish cheeses,

house chutney, celery, frozen grapes, biscuits 7.25



Chef's dessert of the day 5.95

Ask your server for details

Journal

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Appendix B: Menu Questionnaires

Control Questionnaire

Please take a few moments to fill out the following questions...

1. How familiar are you with nutritional labelling for foods?

Very familiar Somewhat familiar Neither familiar nor unfamiliar Somewhat unfamiliar Very unfamiliar

2. Would you find nutritional information (e.g., calories, fat, sugar, salt) useful when choosing your meal?

Very much agree Somewhat agree Neither agree nor disagree Somewhat disagree Very much disagree

3. Would this information influence what you choose from a menu?

Very much agree Somewhat agree Neither agree nor disagree Somewhat disagree Very much disagree

4. What is your gender?

Male Female Prefer not to say

5. Please briefly indicate which dishes you ordered (leave blank if N/A).

Starter	
Main	
Dessert	

Thank you!

Journal

Treatment Questionnaire

Please take a few moments to fill out this short questionnaire...

1. How familiar were you with nutritional labelling before coming to the restaurant?

Very familiar	Somewhat familiar	Neither familiar nor unfamiliar	Somewhat unfamiliar	Very unfamiliar
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. Nutritional labelling influenced what I ordered from the menu.

Very much agree	Somewhat agree	Neither agree nor disagree	Somewhat disagree	Very much disagree
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. Which information on labels did you find **MOST** useful when ordering your meal? (*tick ONE*)

Calories	Walking mins	Fat	Saturated fats	Sugars	Salt
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

4. Which information on labels did you find **LEAST** useful when ordering your meal? (*tick ONE*)

Calories	Walking mins	Fat	Saturated fats	Sugars	Salt
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. What is your gender?

Male Female Prefer not to say

6. Please briefly indicate which dishes you ordered (leave blank if not ordered).

Starter	
Main	
Dessert	

Thank you!

Food Journal